

BELARUS NATIONAL ACADEMY OF SCIENCES  
B.I. STEPANOV's INSTITUTE OF PHYSICS

V.V. Kisel, N.G. Tokarevskaya, A.A. Bogush, V.M. Red'kov<sup>1</sup>

SPIN 1 PARTICLE IN A 15-COMPONENT FORMALISM,  
INTERACTION WITH ELECTROMAGNETIC AND GRAVITATIONAL FIELDS

**Abstract**

A generalized vector particle theory with the use of an extended set of Lorentz group irreducible representations, including scalar, two 4-vectors, and antisymmetric 2-rang tensor, is investigated both in tensor and in matrix approaches. Initial equations depend upon four complex parameters  $\lambda_i$ , obeying two supplementary conditions, so restriction of the model to the case of electrically neutral vector particle is not a trivial task. A special basis in the space of 15-component wave functions is found where instead of four  $\lambda_i$  only one real-valued quantity  $\sigma$ , a bilinear combination of  $\lambda_i$ , is presented. This  $\lambda$ -parameter is interpreted as an additional electromagnetic characteristic of a charged vector particle, polarizability. It is shown that in this basis  $C$ -operation is reduced to the complex conjugation only, without any accompanying linear transformation. The form of  $C$ -operation in the initial basis is calculated too. Invariant bilinear form matrix in both bases are found and the Lagrange formulation of the whole theory is given. Explicit expressions of the conserved current vector and of the energy-momentum tensor are established. In presence of external electromagnetic fields, two supplementary field components, scalar and 4-vector, give a non-trivial contribution to the Lagrangian and conserved quantities. Restriction to a massless vector particle is determined.

Extension of the whole theory to the case of Riemannian space-time is accomplished. Two methods of obtaining corresponding generally covariant wave equations are elaborated: of tensor- and of tetrad-based ones. Their equivalence is proved. It is shown that in case of pure curved space-time models without Cartan torsion no specific additional interaction terms because of non-flat geometry arise. The conformal symmetry of a massless generally covariant equation is demonstrated explicitly. A canonical tensor of energy-momentum  $T_{\beta\alpha}$  is constructed, its conservation law happens to involve the Riemann curvature tensor. Within the framework of known ambiguity of any energy-momentum tensor, a new tensor  $\bar{T}_{\beta\alpha}$  is suggested to be used, which obeys a common conservation law.

36 pages, references — 80

**Key words:** spin 1, non-minimal interaction, polarizability, conformal invariance.

**PASC numbers:** 1130, 0230, 0365, 2110H.

---

<sup>1</sup>E-Mail: redkov@dragon.bas-net.by

## Content

### Introduction

1. Generalization of the vector particle theory
2. Special transformations of the wave function
3. Invariant bilinear form
4. On  $C$ -conjugation
5. Equations in a Riemannian space-time, tensor-based approach
6. Covariant tetrad-based formalism
7. Bilinear invariants in a curved space-time
8. Massless equation and conformal invariance.
9. Canonical energy-momentum tensor

### Results

## Introduction

In order to have a comprehensive theory of higher-spin fields, it is necessary to be able to describe interactions. The most important and the best understood is, of course, the electromagnetic one. But the gravitation is very important as well, at least from theoretical viewpoint if not from practical. Generally, a standpoint may be brought forward that we should give much attention to those facts of physics in the Minkowski space-time which allow for extension to a generally covariant physics. Else one point should be taken as of principal importance. Since wave equations with subsidiary conditions usually lead to consistency difficulties when minimally coupled to an electromagnetic field, it seems best to avoid subsidiary conditions from the start. Furthermore, it seems the best to start with a first-order system, which is linked up to a certain Lagrange function, and in which some physically required restrictive conditions are built in from the very beginning in accordance with the Pauli-Fierz approach [1-2]. The more so is in the context of possible background of a non-Euclidean geometry, in view of arising additional subtleties in a consistency problem.

It is well known that if one looks for a first-order differential equation describing a mass  $m$  spin  $s$  field, that is form-invariant under Lorentz transformations, derivable from a Lagrangian, then one does not uniquely obtain the common Dirac or Duffin-Kemmer or some other equations. A general theory of such first-order and Lagrangian-based equations has been treated at great length Fierz and Pauli [1-2], Dirac [3], Bhabha [4], Harish-Chandra [5], Umezawa [6], and by many others [7-12].


In fact, it was shown that almost an infinite many of such equations is possible. However, there has been a notable dearth of examples of such theories that have been developed at a large extent comparable to the Dirac or Duffin-Kemmer examples. But such particle models, being elaborated in full detail, might shed new light on the theory of general arbitrary-spin wave equations. Much work in this direction has been done [13-39]. For instance, generalized wave equations for particles of spin 1/2 and 1 respectively and with arbitrary anomalous magnetic momentums have been worked out [15,17-19,21,22,24], [28,30,34,37-39]. Else one extension, less known, has been done for boson particle: a next simplest theory beyond the Duffin-Kemmer case was worked out in [20,25,26,35,36] and an explicit representation of the basic first-order equation was given.

In this paper we examine the properties of this new 15-component spin 1 theory in the case of no interaction as well as in presence of external electromagnetic and gravitational fields. It turns out that

this theory is very simply related to the Duffin-Kemmer equation (in recent years some interest in the Duffin-Kemmer-Petiau formalism again can be noted – see [40-45]). In the free-field case, the two theory are equivalent, and the more complicated 15-component model can be reduced to the ordinary 10-component one. In the presence of external electromagnetic fields, one again obtains a Duffin-Kemmer like theory with an additional term corresponding to an electromagnetic polarizability. Thus, this extended theory is definitely inequivalent to the Duffin-Kemmer theory in presence of the minimal electromagnetic coupling. This else one time illustrates explicitly a point emphasized in the literature that although two field theories may be equivalent in absence of interaction, they may be completely inequivalent in presence of interaction.

## 1. Generalization of the vector particle theory

The full physical content of an extended theory of a spin 1 particle will emerge later. As for now, we just postulate certain original set of the Lorentz group representation along with a wave equation for the generalized fields [20,25,26]. Initial generalized equation for a massive spin 1 particle, being associated with the scheme of links



has the form<sup>2</sup> ( $m = Mc/\hbar$ )

$$\begin{aligned} \partial^a(\lambda_1\Phi_{1a} + \lambda_2\Phi_{2a}) - \Phi m &= 0, \\ \mp\lambda_1^*\partial_a\Phi + \lambda_3\partial^b\Phi_{ab} - m\Phi_{1a} &= 0, \\ \pm\lambda_2^*\partial_a\Phi + \lambda_4\partial^b\Phi_{ab} - m\Phi_{2a} &= 0, \\ \partial_a(\lambda_3^*\Phi_{1b} - \lambda_4^*\Phi_{2b}) - \partial_b(\lambda_3^*\Phi_{1a} - \lambda_4^*\Phi_{2a}) - m\Phi_{ab} &= 0. \end{aligned} \quad (1.1)$$

Here all the fields ( $\Phi, \Phi_{1a}, \Phi_{2a}, \Phi_{ab}$ ) as well as  $\lambda_i$  are complex-valued. Therefore restriction of the theory to the case of chargeless particle, which cannot interact with external electromagnetic fields, is to be a matter of special consideration.

It must be noted that the  $\lambda_i$ -s are not fixed uniquely. Any set of such  $\lambda_i$  must obey two restricting conditions [25,26]

$$\lambda_1\lambda_1^* - \lambda_2\lambda_2^* = 0, \quad \lambda_3\lambda_3^* - \lambda_4\lambda_4^* = 1. \quad (1.2)$$

In accordance with eqs. (1.2), freedom of choice of the  $\lambda_i$ -s is as follows:

$$\begin{pmatrix} \lambda'_1 \\ \lambda'_2 \end{pmatrix} = \begin{pmatrix} \cosh\theta & \sinh\theta \\ \sinh\theta & \cosh\theta \end{pmatrix} \begin{pmatrix} \lambda_1 \\ \lambda_2 \end{pmatrix}, \quad \begin{pmatrix} \lambda'_3 \\ \lambda'_4 \end{pmatrix} = \begin{pmatrix} \cosh\Theta & \sinh\Theta \\ \sinh\Theta & \cosh\Theta \end{pmatrix} \begin{pmatrix} \lambda_3 \\ \lambda_4 \end{pmatrix}. \quad (1.3)$$

Generally speaking,  $\theta$ - and  $\Theta$ -transformations are independent of each other.

The tensor equations (1.1) can be formulated as one matrix equation. To this end, let us rewrite eqs. (1.1) in the form

$$\begin{aligned} \partial_a(\lambda_1 g^{ab})\Phi_{1b} + \partial_a(\lambda_2 g^{ab})\Phi_{2b} - m\Phi &= 0, \\ \partial_a(\mp\lambda_1^*\delta_k^a)\Phi - \partial_a(\lambda_3 g^{al}\frac{1}{2}\delta_{lk}^{mn})\Phi_{mn} - m\Phi_{1k} &= 0, \\ \partial_a(\pm\lambda_2^*\delta_k^a)\Phi - \partial_a(\lambda_4 g^{al}\delta_{lk}^{mn})\Phi_{mn} - m\Phi_{2k} &= 0, \\ \partial_a(\lambda_3^*\delta_{kl}^{ab})\Phi_{1b} + \partial_a(-\lambda_4^*\delta_{kl}^{ab})\Phi_{2b} - m\Phi_{kl} &= 0. \end{aligned} \quad (1.4)$$

---

<sup>2</sup>Latin indices take the values 0, 1, 2, 3; at the work in the Minkowski space-time the conventional metric tensor ( $g_{ab}$ ) =  $diag(+1, -1, -1, -1)$ ;  $x_0 = ct$  will be used.

In eqs. (1.4) and in the following a special Kronecker's symbol  $\delta_{kl}^{ab}$ , acting in the space of antisymmetric tensors as a unite operator

$$\delta_{kl}^{ab} = (\delta_k^a \delta_l^b - \delta_l^a \delta_k^b), \quad \frac{1}{2} \delta_{kl}^{ab} \Phi_{ab} = \Phi_{kl}.$$

In turn, the equations (1.4) can be written as a matrix equation:

$$\partial_a \begin{pmatrix} 0 & \lambda_1 g^{ab} & \lambda_2 g^{ab} & 0 \\ \mp \lambda_1^* \delta_k^a & 0 & 0 & -\lambda_3 g^{al} \delta_{lk}^{mn} \\ \pm \lambda_2^* \delta_k^a & 0 & 0 & -\lambda_4 g^{al} \delta_{lk}^{mn} \\ 0 & +\lambda_3^* \delta_{kl}^{ab} & -\lambda_4^* \delta_{kl}^{ab} & 0 \end{pmatrix} \begin{pmatrix} \Phi \\ \Phi_{1b} \\ \Phi_{2b} \\ \Phi_{mn} \end{pmatrix} = m \begin{pmatrix} \Phi \\ \Phi_{1k} \\ \Phi_{2k} \\ \Phi_{kl} \end{pmatrix}, \quad (1.5a)$$

or in a compact form

$$(\Gamma^a \partial_a - m) \Psi = 0, \quad (1.5b)$$

where four matrices  $\Gamma^a$  with the block structure

$$\gamma^a = \begin{pmatrix} 1 & 1 \times 4 & 1 \times 4 & 1 \times 6 \\ 4 \times 1 & 4 \times 4 & 4 \times 4 & 4 \times 6 \\ 4 \times 1 & 4 \times 4 & 4 \times 4 & 4 \times 6 \\ 6 \times 1 & 6 \times 4 & 6 \times 4 & 6 \times 6 \end{pmatrix} \quad (1.5c)$$

have been used. Take notice on the absence of a factor 1/2 in second row of the  $\Gamma^a$  at  $\lambda_3$ ; this is so because we have supposed that  $\Psi$ -function contains only 6 components of the antisymmetric tensor  $\Phi_{kl}$  (those are (01, 02, 03, 23, 31, 12)); so one can sum over 6-combinations  $[kl]$  or can sum over indices  $k$  and  $l$  independently but simultaneously inserting the factor 1/2 (see (1.4)).

The freedom of choice (1.3) can be understood in the following way: let us simultaneously with the change  $\lambda_i \Rightarrow \lambda'_i$  (1.3) perform non-degenerate linear transformation upon  $\Psi$ -function:

$$\Psi' = S \Psi, \quad S = S(\theta, \Theta),$$

$$(\Gamma'^a \partial_a - m) \Psi' = 0, \quad \Gamma'^a = S \Gamma^a S^{-1}, \quad (1.6a)$$

Solving the equation  $\Gamma'^a = S \Gamma^a S^{-1}$ , where  $\Gamma^a$  and  $\Gamma'^a$  are functions of  $\lambda_i$  and  $\lambda'_i$  respectively (see (1.3)), one can easily establish the explicit form of  $S$ :

$$\begin{pmatrix} \Phi'_1 \\ \Phi'_2 \end{pmatrix} = \begin{pmatrix} \cosh \theta & -\sinh \theta \\ -\sinh \theta & \cosh \theta \end{pmatrix} \begin{pmatrix} \Phi_1 \\ \Phi_2 \end{pmatrix}, \quad \Theta = -\theta. \quad (1.6b)$$

Now we are coming to another question. Let us perform a special linear transformation

$$\begin{pmatrix} C_a \\ \Phi_a \end{pmatrix} = \begin{pmatrix} \lambda_1 & \lambda_2 \\ \lambda_3^* & -\lambda_4^* \end{pmatrix} \begin{pmatrix} \Phi_{1a} \\ \Phi_{2a} \end{pmatrix},$$

$$\begin{pmatrix} \Phi_{1a} \\ \Phi_{2a} \end{pmatrix} = \frac{1}{\lambda_1 \lambda_4^* + \lambda_2 \lambda_3^*} \begin{pmatrix} \lambda_4^* & \lambda_2 \\ \lambda_3^* & -\lambda_1 \end{pmatrix} \begin{pmatrix} C_a \\ \Phi_a \end{pmatrix}. \quad (1.7)$$

In these variables, eqs. (1.1) will take on the simpler form (in the following will be convenient instead the scalar function  $\Phi(x)$  to use the notation  $C(x)$ ; so two supplementary components will be  $C(x)$  and  $C_a$ )

$$\begin{aligned} \partial^a C_a - m C &= 0, \\ d \partial^a \Phi_{ba} - m C_b &= 0, \\ \mp d^* \partial_b C + \partial^a \Phi_{ba} - m \Phi_b &= 0, \\ \partial_a \Phi_b - \partial_b \Phi_a - m \Phi_{ab} &= 0, \end{aligned} \quad (1.8)$$

where symbol  $d$  stands for the combination of  $\lambda_i$ -s:  $d = \lambda_1 \lambda_3 + \lambda_2 \lambda_4$ . Take notice that the new system (1.8) depends on  $\lambda_i$  only through  $d$  and  $d^*$ .

In addition it is readily verified that the introduced vectors  $C(x)$  and  $C_a(x)$  are  $\theta$ -invariant. In other words, irrespective of a concrete choice of an initial basis, fixed by a set of  $\lambda_i$ , the final system (1.8) will be the same.

Eqs. (1.8) can be written, as well as eqs. (1.1), in a matrix form:

$$(\Gamma^a \partial_a - m) \Psi = 0 ,$$

$$\partial_a \begin{pmatrix} 0 & g^{ak} & 0 & 0 \\ 0 & 0 & 0 & -dg^{ab} \delta_{bl}^{mn} \\ \mp d^* \delta_l^a & 0 & 0 & -g^{ab} \delta_{bl}^{mn} \\ 0 & 0 & \delta_{bl}^{ak} & 0 \end{pmatrix} \begin{pmatrix} C \\ C_k \\ \Phi_k \\ \Phi_{mn} \end{pmatrix} = m \begin{pmatrix} C \\ C_l \\ \Phi_l \\ \Phi_{bl} \end{pmatrix} . \quad (1.9)$$

Having used two first equations in (1.8) one can exclude the supplementary field  $C(x)$  from third equation in (1.8):

$$\mp \frac{d^* d}{m^2} \partial_a (\partial^k \partial^l \Phi_{kl}) + \partial^b \Phi_{ab} - m \Phi_a = 0 ,$$

$$\partial_a \Phi_b - \partial_b \Phi_a - m \Phi_{ab} = 0 . \quad (1.10a)$$

As the term  $(\partial^k \partial^l \Phi_{kl})$  vanishes identically. Thus, in free case (absence of external fields) additional parameter of the extended theory of the spin 1 particle does not manifest itself anyhow. At this two supplementary field components are as follows:

$$C = 0 , \partial^a C_a = 0 , C_a = d \Phi_a , \quad (1.10b)$$

and the main equations are

$$\partial^a \Phi_{ba} - m \Phi_b = 0 , \partial_a \Phi_b - \partial_b \Phi_a - m \Phi_{ab} = 0 . \quad (1.11)$$

these are ordinary Proca equations.

Other situation is realized in presence of external electromagnetic fields. Making conventional change in eqs. (1.8)

$$\partial_a \implies D_a = \partial_a - ig A_a(x) ,$$

$g = \frac{e}{\hbar c}$ ,  $e$  — electric charge of the particle, instead of (1.8) we get

$$D^a C_a - m C = 0 , d D^a \Phi_{ba} - m C_b = 0 ,$$

$$\mp d^* D_b C + D^a \Phi_{ba} - m \Phi_b = 0 ,$$

$$D_a \Phi_b - D_b \Phi_a - m \Phi_{ab} = 0 . \quad (1.12)$$

From second equation in (1.12) it follows

$$D^a C_a = \frac{d}{m} D^a D^b \Phi_{ab} = i \frac{ed}{m} \frac{1}{2} F^{ab} \Phi_{ab} , \quad (1.13a)$$

where  $F_{ab} = (\partial_a A_b - \partial_b A_a)$ . Accounting for (1.13a), from first equation in (1.12) we get

$$C = -i \frac{ed}{m^2} \frac{1}{2} F^{ab} \Phi_{ab} . \quad (1.13b)$$

Substituting (1.13b) into third equation in (1.12), we arrive at

$$D^a C_a - m C = 0 , d D^a \Phi_{ba} - m C_b = 0 ,$$

$$\pm i \frac{edd^*}{m^2} D_k \left( \frac{1}{2} F^{ab} \Phi_{ab} \right) + D^a \Phi_{ak} - m \Phi_k = 0 ,$$

$$D_a \Phi_b - D_b \Phi_a - m \Phi_{ab} = 0 . \quad (1.14)$$

Equations (1.14) describe behavior of the vector particle in external electromagnetic field. Two first relations in (1.14) are to be considered as additional, they give us possibility to construct supplementary fields  $C(x), C_a$  in terms of the main  $\Phi_a, \Phi_{ab}$ . In (1.14)  $d^* d$ -dependent term is an additional one and it represents some electromagnetic characteristic of the particle.

Now we are coming to another question and will consider how one should determine the massless limit of the theory. The corresponding equations are to be

$$\partial^a C_a - C = 0 , d \partial^a \Phi_{ba} - C_b = 0 ,$$

$$\begin{aligned} \mp d^* \partial_b C + \partial^a \Phi_{ba} &= 0 , \\ \partial_a \Phi_b - \partial_b \Phi_a - \Phi_{ab} &= 0 . \end{aligned} \quad (1.15)$$

This system is invariant under gauge transformation

$$\begin{aligned} \Phi_a(x) &\Longrightarrow \Phi'_a(x) = \Phi_a(x) + \partial_a Z(x) , \\ C(x) &= inv , C_a(x) = inv , \Phi_{ab}(x) = inv . \end{aligned} \quad (1.16)$$

Having all the fields complex, the  $Z(x)$  is to be complex too. It should be borne in mind that such a different behavior of two vector components  $C_a, \Phi_a$  under gauge transformation (see. (1.16) ) holds only in the chosen basis. Taking the gauge transformation (1.16) to the initial basis  $\Phi_{1a}, \Phi_{2a}$  (1.1) we will have

$$\Phi_{1a} \Longrightarrow \Phi_{1a} + \frac{\lambda_1^*}{d^*} \partial_a \Lambda(x) , \quad \Phi_{2a} \Longrightarrow \Phi_{2a} - \frac{\lambda_2^*}{d^*} \partial_a \Lambda(x) , \quad (1.17)$$

both vectors transform simultaneously.

In the massless case the supplementary fields turn out to be zero

$$C = 0 , C_b = 0 ,$$

Thus, eq. (1.15) takes on the form of the ordinary Proca massless system

$$\partial^a \Phi_{ba} = 0 , \partial_a \Phi_b - \partial_b \Phi_a - \Phi_{ab} = 0 . \quad (1.18)$$

This is quite understandable result, having remembered that in massive case the  $\lambda_i$  exhibit themselves physically only in presence of external electromagnetic fields  $A_a(x)$ .

However, from heuristic considerations (having in mind the gauge invariance principle), in the massless case as well one can theoretical possibility, to study a massless complex-valued field in external vector ('electromagnetic') field<sup>3</sup> At this eqs. (1.15) will change into

$$\begin{aligned} D^a C_a - C &= 0 , \\ d D^a \Phi_{ba} - C_b &= 0 , \\ \mp d^* D_b C + D^a \Phi_{ba} &= 0 , \\ D_a \Phi_b - D_b \Phi_a - \Phi_{ab} &= 0 , \end{aligned} \quad (1.19)$$

where  $D_a = \partial_a - ig A_a$ . Eqs. (1.19) are invariant under  $U(1)$ -gauge transformation, different from previous gauge symmetry (1.16):

$$A_a \Longrightarrow A_a + \partial_a \Lambda(x) , (C, C_a, \Phi_a, \Phi_{ab}) \Longrightarrow e^{ig\Lambda} (C, C_a, \Phi_a, \Phi_{ab}) . \quad (1.20)$$

It can be readily verified that eqs. (1.19) are equivalent to the following ones

$$\begin{aligned} C_b &= d D^a \Phi_{ba} , \\ C &= -i ed \frac{1}{2} F^{ab} \Phi_{ab} , \\ \pm i ed^* d D_b (\frac{1}{2} F^{kl} \Phi_{kl}) + D^a \Phi_{ab} &= 0 , \\ D_a \Phi_b - D_b \Phi_a - \Phi_{ab} &= 0 . \end{aligned} \quad (1.21)$$

So, in a massless case, in presence of external vector fields  $A_a$ , the  $\lambda_i$ -parameters manifest themselves as well.

---

<sup>3</sup>The term 'electromagnetic' here should be understood with caution, in fact as a matter of convention.

## 2. Special transformations of the wave function

In Sec. 2 we are going to give special attention to the question of the charge-symmetry. At this, for simplicity, instead of  $D_a = \partial_a - igA_a$  we will write the operator  $\partial_a$ , but remembering that everything concerns the charged particle as well (with taking  $g$  into  $-g$  at  $C$ -conjugation).

Let us turn to eqs. (1.8) and write down complex-conjugate ones:

$$\begin{aligned} \partial^a C_a^* - m C^* &= 0, \\ d^* \partial^a \Phi_{ba}^* - m C_b^* &= 0, \\ \mp d \partial_b C^* + \partial^a \Phi_{ba}^* - m \Phi_b^* &= 0, \\ \partial_a \Phi_b^* - \partial_b \Phi_a^* - m \Phi_{ab}^* &= 0. \end{aligned} \quad (2.1)$$

Defining  $C$ -conjugate wave functions according to

$$\begin{aligned} (d/d^*) C^* &= C^{(c)}, \quad (d/d^*) C_a^* = C_a^{(c)}, \\ \Phi_a^* &= \Phi_a^{(c)}, \quad \Phi_{ab}^* = \Phi_{ab}^{(c)}, \end{aligned} \quad (2.2)$$

eqs. (2.1) can be rewritten as

$$\begin{aligned} \partial^a C_a^{(c)} - m C^{(c)} &= 0, \quad d \partial^a \Phi_{ba}^{(c)} - m C_b^{(c)} = 0, \\ \mp d^* \partial_b C^{(c)} + \partial^a \Phi_{ba}^{(c)} - m \Phi_b^{(c)} &= 0, \\ \partial_a \Phi_b^{(c)} - \partial_b \Phi_a^{(c)} - m \Phi_{ab}^{(c)} &= 0, \end{aligned}$$

which exactly coincides with eqs. (1.8). Therefore eqs. (1.8) are  $C$ -invariant and  $C$ -conjugation matrix has the form

$$\Psi^c = C \Psi^*, \quad C = \begin{pmatrix} (d/d^*) & 0 & 0 & 0 \\ 0 & (d/d^*) I_4 & 0 & 0 \\ 0 & 0 & I_4 & 0 \\ 0 & 0 & 0 & I_6 \end{pmatrix}. \quad (2.3)$$

Now, let us show that there exists a basis in which  $S$ -operation has a simpler form. To this end, turning again to eqs. (1.8), let us separate out phase factors from  $d$  and  $d^*$ :  $d \implies d_0 e^{it}$ ,  $d^* \implies d_0 e^{-it}$ . Then eqs. (1.8) will read as

$$\begin{aligned} \partial^a C_a - m C &= 0, \\ d_0 e^{+it} \partial^a \Phi_{ba} - m C_b &= 0, \\ \mp d_0 e^{-it} \partial_b C + \partial^a \Phi_{ba} - m \Phi_b &= 0, \\ \partial_a \Phi_b - \partial_b \Phi_a - m \Phi_{ab} &= 0. \end{aligned} \quad (2.4)$$

Now, defining new fields

$$e^{-it} C(x) \implies C(x), \quad e^{-it} C_a(x) \implies C_a(x), \quad (2.5)$$

we take eqs. (2.4) into

$$\begin{aligned} \partial^a C_a - m C &= 0, \\ d_0 \partial^a \Phi_{ba} - m C_b &= 0, \\ \mp d_0 \partial_b C + \partial^a \Phi_{ba} - m \Phi_b &= 0, \\ \partial_a \Phi_b - \partial_b \Phi_a - m \Phi_{ab} &= 0, \end{aligned} \quad (2.6)$$

here  $d_0$  is a real-valued quantity.

Matrix form of eqs. (2.6) is

$$\partial_a \begin{pmatrix} 0 & g^{ak} & 0 & 0 \\ 0 & 0 & 0 & -d_0 g^{ab} \delta_{bl}^{mn} \\ \mp d_0 \delta_l^a & 0 & 0 & -g^{ab} \delta_{bl}^{mn} \\ 0 & 0 & \delta_{bl}^{ak} & 0 \end{pmatrix} \begin{pmatrix} C \\ C_k \\ \Phi_k \\ \Phi_{mn} \end{pmatrix} = m \begin{pmatrix} C \\ C_l \\ \Phi_l \\ \Phi_{bl} \end{pmatrix}. \quad (2.7)$$

Evidently, in this basis  $C$ -operation is reduced to the complex conjugation only:

$$\Psi^c(x) = \Psi^*(x) . \quad (2.8)$$

Else one transformation upon  $\Psi$  is to be done:

$$d_0^{-1} C(x) \Longrightarrow C'(x) , \quad d_0^{-1} C_a(x) \Longrightarrow C'_a(x) , \quad (2.9)$$

then eqs. (2.6) read as

$$\begin{aligned} \partial^a C_a - mC &= 0 , \\ \partial^a \Phi_{ba} - mC_b &= 0 , \\ \mp d_0^2 \partial_b C + \partial^a \Phi_{ba} - m\Phi_b &= 0 , \\ \partial_a \Phi_b - \partial_b \Phi_a - m\Phi_{ab} &= 0 . \end{aligned} \quad (2.10)$$

In the following this  $d_0^2$  will be designated as  $\sigma$ . Also it is quite understandable that existence of two possibilities in eqs. (2.10), associated with two signs  $\pm$ , can be described as different in sign quantities  $\sigma$ . In other words, these signs  $\pm$  at  $\sigma$  can be omitted. It should be mentioned that  $\sigma$ -parameter is dimensionless whereas all the component of  $\Psi$  have dimension ( $l = cm$ :  $[C] = [C_a] = [\Phi_a] = [\Phi_{ab}] = l^{-3/2}$ ). Matrix form of eqs. (2.10) is as follows

$$\partial_a \begin{pmatrix} 0 & g^{ak} & 0 & 0 \\ 0 & 0 & 0 & -g^{ab}\delta_{bl}^{mn} \\ \sigma\delta_l^a & 0 & 0 & -g^{ab}\delta_{bl}^{mn} \\ 0 & 0 & \delta_{bl}^{ak} & 0 \end{pmatrix} \begin{pmatrix} C \\ C_k \\ \Phi_k \\ \Phi_{mn} \end{pmatrix} = m \begin{pmatrix} C \\ C_l \\ \Phi_l \\ \Phi_{bl} \end{pmatrix} . \quad (2.12)$$

In massless limit, instead of (2.11) we will have

$$\begin{aligned} \partial^a C_a - C &= 0 , \\ \partial^a \Phi_{ba} - C_b &= 0 , \\ \sigma \partial_b C + \partial^a \Phi_{ba} &= 0 , \\ \partial_a \Phi_b - \partial_b \Phi_a - \Phi_{ab} &= 0 . \end{aligned} \quad (2.13)$$

To avoid misunderstanding it should be noted that though the same symbol  $\sigma$  is used both in (2.11) and (2.13) but this symbol stands for quite different characteristics. In massless case  $\sigma$  must have dimension of squared length.

### 3. Invariant bilinear form

Let us consider the question about explicit expression of invariant bilinear form matrix in this theory. In the work with the equation (see (2.12))

$$(\Gamma^a \partial_a - m) \Psi = 0 \quad (3.1)$$

we will employ the block structure of the  $\Gamma^a$ -s

$$\Gamma^a = \begin{pmatrix} 0 & G^a & 0 & 0 \\ 0 & 0 & 0 & K^a \\ \sigma\Delta^a & 0 & 0 & K^a \\ 0 & 0 & \Lambda^a & 0 \end{pmatrix} , \quad (3.2a)$$

where

$$\begin{aligned} (K^a)_l{}^{mn} &= (-g^{am}\delta_l^n + g^{an}\delta_l^m) , \\ (\Lambda^a)_{bl}{}^k &= \delta_{bl}^{ak} , \\ (G^a)_{(0)}{}^k &= g^{ak} , \\ (\Delta^a)_l{}^{(0)} &= \delta_l^a . \end{aligned} \quad (3.2b)$$



As known, required matrix  $\eta$  of invariant form is determined by the relation [12]

$$\eta^{-1}(\Gamma^a)^+\eta = -\Gamma^a. \quad (3.3)$$

If the  $\eta$  is found, then  $\bar{\Psi} = \Psi^+\eta$  will obey

$$\bar{\Psi} (\Gamma^a \overleftarrow{\partial}_a + m) = 0. \quad (3.4)$$

Allowing for (3.1) and (3.4), one can straightforwardly obtain this conventional expression for a current conserved:

$$J^a = \bar{\Psi}\Gamma^a\Psi, \partial_a J^a = 0. \quad (3.4)$$

By definition, a Lorentz invariant is to be constructed with the use of  $\eta$  according to formula  $\Psi^+\eta\Psi$ . Its most general structure is as follows

$$\Psi^+\eta\Psi = r_1 C^*C + r_2 C_a^*C^a + r_3 C_a^*\Phi^a + r_4 \Phi_a^*C^a + r_5 \Phi_a^*\Phi^a + \frac{1}{2}r_6 \Phi_{ab}^*\Phi_{ab},$$

and hence  $\eta$  can be as

$$\eta = \begin{pmatrix} R_1 & 0 & 0 & 0 \\ 0 & R_2 & R_3 & 0 \\ 0 & R_4 & R_5 & 0 \\ 0 & 0 & 0 & R_6 \end{pmatrix} = \begin{pmatrix} r_1 & 0 & 0 & 0 \\ 0 & r_2 g^{ab} & r_3 g^{ab} & 0 \\ 0 & r_4 g^{ab} & r_5 g^{ab} & 0 \\ 0 & 0 & 0 & r_6 H^{mn,ab} \end{pmatrix}. \quad (3.5)$$

Here  $H^{mn,ab} = (g^{ma}g^{nb} - g^{mb}g^{na})$ . Taking into account representation of  $\Gamma^a, (\Gamma^a)^+$  and  $\eta$  in terms of block-matrices, from eq. (3.3) rewritten as  $\Gamma^{a+}\eta + \eta\Gamma^a = 0$ , we readily get to the set of relations

$$\begin{aligned} \sigma\tilde{\Delta}^a R_4 + R_1 G^a &= 0, \quad R_5 = 0, \\ \tilde{G}^a R_1 + \sigma R_3 \Delta^a, \quad R_2 + R_3 &= 0, \\ R_5 = 0, \quad \tilde{\Lambda}^a R_6 + R_4 K^a &= 0, \\ R_2 + R_4 = 0, \quad \tilde{K}^a R_3 + R_6 \Lambda^a &= 0. \end{aligned} \quad (3.6a)$$

In (3.6) one can distinguish couples of equations transformed into each other on matrix transposing; therefore each couple provides us with a single equation. Allowing for formulas (3.2), from eqs. (3.6) it follows

$$r_1 = +\sigma r_2, \quad r_3 = -r_2, \quad r_4 = -r_2, \quad r_5 = 0, \quad r_6 = -r_2. \quad (3.6b)$$

Thus, in the used basis  $\eta$ -matrix is given by

$$\eta = \text{const} \begin{pmatrix} +\sigma & 0 & 0 & 0 \\ 0 & +g^{ab} & -g^{ab} & 0 \\ 0 & -g^{ab} & 0 & 0 \\ 0 & 0 & 0 & -H^{mn,ab} \end{pmatrix}. \quad (3.7)$$

The relevant feature of  $\eta$ -matrix is that the invariant-form matrix has turned out to be of non-diagonal structure.

But this property does not hold in all other possible bases. For instance, let us show that in a representation associated with eqs. (1.1) the  $\eta$ -matrix will have a common diagonal form. Actually, being linked up to eqs. (1.8), the  $\Gamma^a$ -s are

$$\Gamma^a = \begin{pmatrix} 0 & \lambda_1 G^a & \lambda_2 G^a & 0 \\ -\nu\lambda_1^* \Delta^a & 0 & 0 & \lambda_3 K^a \\ +\nu\lambda_2^* \Delta^a & 0 & 0 & \lambda_4 K^a \\ 0 & \lambda_3^* \Lambda^a & -\lambda_4^* \Lambda^a & 0 \end{pmatrix}, \quad (3.8)$$

here the same notation for block-matrices  $G^a, \Delta^a, K^a, \Lambda^a$  as in (3.2b)  $\eta$  has been used. In contrast to the above, here an appropriate substitution for  $\eta$ -matrix happens to be of typical diagonal form

$$\eta = \begin{pmatrix} C_1 & 0 & 0 & 0 \\ 0 & C_2 & 0 & 0 \\ 0 & 0 & C_3 & 0 \\ 0 & 0 & 0 & C_4 \end{pmatrix} = \begin{pmatrix} c_1 & 0 & 0 & 0 \\ 0 & c_2 g^{ab} & 0 & 0 \\ 0 & 0 & c_3 g^{ab} & 0 \\ 0 & 0 & 0 & c_4 H^{ab,mn} \end{pmatrix}. \quad (3.9)$$

Correspondingly, the relation  $(\Gamma^a)^+ \eta + \eta \Gamma^a = 0$  yields

$$\begin{aligned} -\nu \tilde{\Delta}^a C_2 + C_1 G^a &= 0, \quad \nu \tilde{\Delta}^a C_3 + C_1 G^a = 0, \\ \tilde{G}^a C_1 - \nu C_2 \Delta^a, \quad \tilde{\Lambda}^a C_4 + C_2 K^a, \\ \tilde{G}^a C_1 + \nu C_3 \Delta^a &= 0, \quad -\tilde{\Lambda}^a C_4 + C_3 K^a = 0, \\ \tilde{K}^a C_2 + C_4 \Lambda^a &= 0, \quad \tilde{K}^a C_3 - C_4 \Lambda^a = 0, \end{aligned}$$

from which one can easily derive

$$c_2 = +c_1 \nu, \quad c_3 = -c_1 \nu, \quad c_4 = +c_1 \nu.$$

Therefore, we have got to

$$\eta = c \begin{pmatrix} -\nu & 0 & 0 & 0 \\ 0 & -g^{ab} & 0 & 0 \\ 0 & 0 & +g^{ab} & 0 \\ 0 & 0 & 0 & -H^{ab,mn} \end{pmatrix}. \quad (3.10)$$

One can give else one (additional) treatment of the matter: namely, to follow how the expression for  $\eta$  in the final basis (3.7) might be calculated on translating that of the initial  $\eta$  according to (3.10). The rule for transforming an invariant-form matrix at changing bases in  $\Psi$ -space can be derived quite easily. From relations defining  $\eta$ :

$$\bar{\Psi} = \Psi^+ \eta, \quad \eta^{-1} \Gamma^{a+} \eta = -\Gamma^a,$$

will immediately follow analogous ones for defining  $\eta'$

$$\bar{\Psi}' = \Psi'^+ \eta', \quad (\eta')^{-1} \Gamma'^{a+} \eta' = -\Gamma'^a,$$

if the following relationships

$$\eta' = (S^+)^{-1} \eta S^{-1}, \quad \text{or} \quad \eta = (S^+) \eta' S. \quad (3.11)$$

hold.

With the use of eq. (3.11) let us transform the  $\eta$ -matrix (3.10) to a new basis. At this it will be convenient to take in mind the following scheme (see. Sec 2)

$$\begin{array}{ccccccc} \lambda_i & \xrightarrow{S_1} & \{d, d^*\} & \xrightarrow{S_2} & \{|d|, |d|\} & \xrightarrow{S_3} & \{\sigma = |d|^2\} \\ \Psi, \eta & & \Psi', \eta' & & \Psi'', \eta'' & & \Psi''', \eta''' \end{array}$$

First, let us perform the transformation  $\Psi' = S_1 \Psi$ . It will be more convenient to employ the formula  $S_1^+ \eta' S_1 = \eta$ , where

$$S_1 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \lambda_1 & \lambda_2 & 0 \\ 0 & \lambda_3^* & -\lambda_4^* & 0 \\ 0 & 0 & 0 & I_6 \end{pmatrix}, \quad S_1^+ = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \lambda_1^* & \lambda_3 & 0 \\ 0 & \lambda_2^* & -\lambda_4 & 0 \\ 0 & 0 & 0 & I_6 \end{pmatrix}. \quad (3.12)$$

Thus, allowing for the substitution for  $\eta'$

$$\eta' = \begin{pmatrix} R_1 & 0 & 0 & 0 \\ 0 & R_2 & R_3 & 0 \\ 0 & R_4 & R_5 & 0 \\ 0 & 0 & 0 & R_6 \end{pmatrix},$$

with the use of the relations  $S_1^+ \eta' S_1 = \eta$  and (see. (3.10))  $r_1 = -c\nu$ ,  $R_6 = c H$ , one can derive the following

$$(\lambda_1^* R_2 + \lambda_3 R_4) \lambda_1 + (\lambda_1^* R_3 + \lambda_3 R_5) \lambda_3^* = -c,$$

$$\begin{aligned}
(\lambda_1^* R_2 + \lambda_3 R_4) \lambda_2 - (\lambda_1^* R_3 + \lambda_3 R_5) \lambda_4^* &= 0, \\
(\lambda_2^* R_2 - \lambda_4 R_4) \lambda_1 + (\lambda_2^* R_3 - \lambda_4 R_5) \lambda_3^* &= 0, \\
(\lambda_2^* R_2 - \lambda_4 R_4) \lambda_2 - (\lambda_2^* R_3 - \lambda_4 R_5) \lambda_4^* &= +c.
\end{aligned} \tag{3.13}$$

With the notation

$$\begin{aligned}
\lambda_1^* r_2 + \lambda_3 r_4 &= A, & \lambda_1^* r_3 + \lambda_3 r_5 &= B \\
\lambda_2^* r_2 - \lambda_4 r_4 &= a, & \lambda_2^* r_3 - \lambda_4 r_5 &= b
\end{aligned} \tag{3.14}$$

the system (3.13) reads as

$$\begin{cases} A\lambda_1 + B\lambda_3^* = -c \\ A\lambda_2 - B\lambda_4^* = 0 \end{cases}, \quad \begin{cases} a\lambda_1 + b\lambda_3^* = 0 \\ a\lambda_2 - b\lambda_4^* = +c \end{cases}. \tag{3.15}$$

From (3.15) it follows

$$\begin{aligned}
A &= -\frac{\lambda_4^*}{\lambda_1 \lambda_4^* + \lambda_2 \lambda_3^*} c, & B &= -\frac{\lambda_2}{\lambda_1 \lambda_4^* + \lambda_2 \lambda_3^*} c, \\
a &= +\frac{\lambda_3^*}{\lambda_1 \lambda_4^* + \lambda_2 \lambda_3^*} c, & b &= -\frac{\lambda_1}{\lambda_1 \lambda_4^* + \lambda_2 \lambda_3^*} c,
\end{aligned}$$

In turn, from (3.14) we obtain

$$\begin{aligned}
r_2 &= \frac{A\lambda_4 + a\lambda_3}{\lambda_1^* \lambda_4 + \lambda_2^* \lambda_3}, & r_4 &= \frac{A\lambda_2^* - a\lambda_1^*}{\lambda_1^* \lambda_4 + \lambda_2^* \lambda_3}, \\
r_3 &= \frac{B\lambda_4 + b\lambda_3}{\lambda_1^* \lambda_4 + \lambda_2^* \lambda_3}, & r_5 &= \frac{B\lambda_2^* - b\lambda_1^*}{\lambda_1^* \lambda_4 + \lambda_2^* \lambda_3}.
\end{aligned}$$

From this, accounting for the above expressions for  $A, B, a, b$ , we get to

$$r_2 = \frac{+c}{DD^*}, \quad r_4 = \frac{-c}{DD^*} d^*, \quad r_3 = \frac{-c}{DD^*} d, \quad r_5 = 0, \tag{3.16}$$

where  $D = \lambda_1^* \lambda_4 + \lambda_2^* \lambda_3$ . With  $\lambda_1 \lambda_1^* = \lambda_2 \lambda_2^*$ , the expression for  $D$  can be rewritten as

$$D = \lambda_1^* \lambda_4 + \lambda_2^* \lambda_3 = \lambda_1^* (\lambda_4 + \frac{\lambda_2^*}{\lambda_1^*} \lambda_3) = \lambda_1^* (\lambda_4 + \frac{\lambda_1}{\lambda_2} \lambda_3) = \frac{\lambda_1}{\lambda_2} d;$$

and correspondingly the equality

$$D D^* = \frac{\lambda_1}{\lambda_2} d \frac{\lambda_1^*}{\lambda_2^*} d^* = d d^*$$

holds. Therefore, we have arrived at the result:

$$r_1 = -c\nu, \quad r_6 = +c, \quad r_2 = +\frac{1}{d^* d} c, \quad r_3 = -\frac{1}{d^*} c, \quad r_4 = -\frac{1}{d} c, \quad r_5 = 0, \tag{3.17a}$$

that is

$$\eta' = c \begin{pmatrix} -\nu & 0 & 0 & 0 \\ 0 & (dd^*)^{-1} g^{ab} & -(d^*)^{-1} g^{ab} & 0 \\ 0 & -d^{-1} g^{ab} & 0 & 0 \\ 0 & 0 & 0 & +H^{ab,mn} \end{pmatrix}. \tag{3.17b}$$

Now let us perform else one transformation

$$\Psi'' = S_2 \Psi', \quad S_2 = \begin{pmatrix} e^{-it} & 0 & 0 & 0 \\ 0 & e^{-it} I_4 & 0 & 0 \\ 0 & 0 & I_4 & 0 \\ 0 & 0 & 0 & I_6 \end{pmatrix}. \tag{3.18a}$$

Correspondingly for  $\eta''$  we will have

$$\eta'' = c \begin{pmatrix} -\nu & 0 & 0 & 0 \\ 0 & d_0^{-2} g^{ab} & -d_0^{-1} g^{ab} & 0 \\ 0 & -d_0^{-1} g^{ab} & 0 & 0 \\ 0 & 0 & 0 & +H^{ab,mn} \end{pmatrix}. \tag{3.18b}$$

And finally, third step:

$$\begin{aligned} C''' &= d_0^{-1} C'' , \quad C_a''' = d_0^{-1} \Phi_a C_a'' , \\ \Phi_a''' &= \Phi_a'' , \quad \Phi_{ab}''' = \Phi_{ab}'' , \end{aligned} \quad (3.19a)$$

and further

$$\eta''' = c \begin{pmatrix} -\nu d_0^2 & 0 & 0 & 0 \\ 0 & +g^{ab} & -g^{ab} & 0 \\ 0 & -g^{ab} & 0 & 0 \\ 0 & 0 & 0 & -H^{ab,mn} \end{pmatrix} . \quad (3.19b)$$

The  $\eta'''$  will coincide with (3.7), if the relation  $-\nu_0 d^2 = \sigma$  is taken into account.

With the use of the expression (3.7) for  $\eta$ , the current (3.5) can be brought to a tensor form:

$$\Psi^+ \eta \Gamma^a \Psi = \text{const} \Psi^+ \begin{pmatrix} +\sigma & 0 & 0 & 0 \\ 0 & +g^{\cdot\cdot} & -g^{\cdot\cdot} & 0 \\ 0 & -g^{\cdot\cdot} & 0 & 0 \\ 0 & 0 & 0 & -H^{\cdot\cdot\cdot\cdot} \end{pmatrix} \Gamma^a \Psi ,$$

and further

$$J^a = \text{const} \{ +\sigma (C^* C^a - C C^{*a}) - (\Phi^{*al} \Phi_l - \Phi^{al} \Phi_l^*) \} . \quad (3.20)$$

Since, in free case, the scalar component  $C(x)$  vanishes (see (1.10)), the current (3.20) will coincide with the ordinary expression for a vector particle current. However, in presence of external electromagnetic fields, the current of generalized theory contains a non-zero additional term proportional to the  $\sigma$ -factor (see (3.20)).

Lagrangian of the theory can be written down straightforwardly in a matrix form

$$L = +\frac{1}{2} \bar{\Psi} (\Gamma^a \vec{\partial}_a - m) \Psi - \frac{1}{2} \bar{\Psi} (\Gamma^a \overleftarrow{\partial}_a + m) \Psi \quad (3.21)$$

with Euler-Lagrange equations as follows

$$\frac{\partial L}{\partial \bar{\Psi}} - \partial_a \frac{\partial L}{\partial \bar{\Psi}_{,a}} = 0 , \quad \frac{\partial L}{\partial \Psi} - \partial_a \frac{\partial L}{\partial \Psi_{,a}} = 0$$

having explicit forms (3.4) and (3.1) respectively. Tensor representation of the Lagrangian (3.21) in basis (2.10) will follow immediately by taking into account the above block structure of all the quantities involved. At this it is useful to have fixed some intermediate steps. Having employed the notation

$$[ C(x), C_l(x), \Phi_l(x), \Phi_{kl}(x) ] = (\Phi, C, A, F) . \quad (3.22)$$

we have

$$\begin{aligned} \bar{\Psi} \Psi &= [ \sigma \Phi^* \Phi + C^* g C - (A^* g C + C^* g A) - F^* H F ] , \\ \bar{\Psi} \Gamma^a \vec{\partial}_a \Psi &= \\ &= [ \sigma (-C^* g \Delta^a \partial_a \Phi + \Phi^* G^a \partial_a C) - F^* H \Lambda^a \partial_a A - A^* g K^a \partial_a F ] , \\ \bar{\Psi} \Gamma^a \overleftarrow{\partial}_a \Psi &= \\ &= [ \sigma (-C^* g \Delta^a \overleftarrow{\partial}_a \Phi + \Phi^* G^a \overleftarrow{\partial}_a C) - F^* H \Lambda^a \overleftarrow{\partial}_a A - A^* g K^a \overleftarrow{\partial}_a F ] , \end{aligned} \quad (3.23a)$$

and further

$$\begin{aligned} \bar{\Psi} \Psi &= [ \sigma C^* C + C_l^* C^l - (\Phi_l^* C^l + \Phi_l C^{*l}) - \frac{1}{2} \Phi_{kl}^* \Phi^{kl} ] , \\ \bar{\Psi} \Gamma^a \vec{\partial}_a \Psi &= [ +\sigma (-C^{*l} \partial_l C + C^* \partial_l C^l) - \Phi^{*lk} \partial_l \Phi_k - \Phi_l^* \partial_k \Phi^{lk} ] , \\ \bar{\Psi} \Gamma^a \overleftarrow{\partial}_a \Psi &= \\ &= [ +\sigma (-C^{*l} \overleftarrow{\partial}_l C + C^* \overleftarrow{\partial}_l C^l) - \Phi^{*lk} \overleftarrow{\partial}_l \Phi_k - \Phi_l^* \overleftarrow{\partial}_k \Phi^{lk} ] , \end{aligned} \quad (3.23b)$$

Therefore, the Lagrangian (3.21) reads explicitly as

$$L = \frac{1}{2} [ -\sigma (C^{*l} \partial_l C + C^l \partial_l C^*) + \sigma (C^* \partial_l C^l + C \partial_l C^{*l}) -$$

$$\begin{aligned}
& - ( \Phi^{*lk} \partial_l \Phi_k + \Phi^{lk} \partial_l \Phi_k^* ) - ( \Phi_l^* \partial_k \Phi^{lk} + \Phi_l \partial_k \Phi^{*lk} ) ] - \\
& - m [ \sigma C^* C + C_l^* C^l - ( \Phi_l^* C^l + \Phi_l C^{*l} ) - \frac{1}{2} \Phi_{kl}^* \Phi^{kl} ] .
\end{aligned} \tag{3.24}$$

One important feature of this representation (3.24) for  $L$  is worth to be should noted specially. The matter is that since in free case two relationships

$$C(x) = 0, \quad C_l(x) = +\Phi_l(x) \tag{3.25}$$

hold, then the Lagrangian (3.24) reduces to the conventional Lagrangian for ordinary vector particle

$$\begin{aligned}
L^0 = \frac{1}{2} [ & - ( \Phi^{*lk} \partial_l \Phi_k + \Phi^{lk} \partial_l \Phi_k^* ) - ( \Phi_l^* \partial_k \Phi^{lk} + \Phi_l \partial_k \Phi^{*lk} ) ] - \\
& - m [ \Phi_l \Phi^{*l} - \frac{1}{2} \Phi_{kl}^* \Phi^{kl} ] ,
\end{aligned} \tag{3.26}$$

To prevent a possible misunderstanding else one thing must be pointed out: going from  $L$  (3.24) to  $L^0$  (3.26) is realized only in the manner just described but by no means through simple imposing  $\sigma = 0$  condition.

And finally, to else one fact should be given a special attention. Because of a non-diagonal structure of the invariant-form matrix  $\eta$  in the basis used, the  $\bar{\Psi}$ -function conjugated has the following explicit form

$$\bar{\Psi}(x) = [ \sigma C^*, (C^{*l} - \Phi^{*l}), -C^{*l}, -\Phi^{*kl} ] , \tag{3.27}$$

second and third equations in (2.11) can be found on variation of Lagrangian (3.24) with respect to  $(C^{*l} - \Phi^{*l})$  and  $(-C^{*l})$  respectively.

Going from the Lagrangian (3.21) to a massless case is achieved by a single formal change

$$m ( \bar{\Psi} \Psi ) \implies \bar{\Psi} P \Psi, \quad P = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & I & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & I \end{pmatrix}, \tag{3.28a}$$

from which it follows (see (3.26))

$$\begin{aligned}
& -m [ \sigma C^* C + C_l^* C^l - ( \Phi_l^* C^l + \Phi_l C^{*l} ) - \frac{1}{2} \Phi_{kl}^* \Phi^{kl} ] \implies \\
& - [ \sigma C^* C + C_l^* C^l - \frac{1}{2} \Phi_{kl}^* \Phi^{kl} ] .
\end{aligned} \tag{3.28b}$$

#### 4. On $C$ -conjugation

As it was shown above, the  $C$ -operation for the generalized field is described as a pure complex conjugation (without any accompanying linear transformation over  $\Psi$ -function) in the basis (2.6). Now we are going to see how will look  $C$ -operation in the initial basis (1.1). It will be useful to consider this question in two different ways: first, solving a defining relation for this operation in the basis (1.1), and alternatively on direct translating the known  $C$ -form in basis (2.6) into initial one.

Let  $C$ -matrix in a  $\Psi(x)$ -basis be known; in other words, the following relationships

$$\begin{aligned}
(\Gamma^a \partial_a - M) \Psi(x) &= 0, & (\Gamma^a \partial_a - M) \Psi^c(x) &= 0, \\
\Psi^c(x) &= C \Psi^*(x), & C(\Gamma^a)^* C^{-1} &= +\Gamma^a
\end{aligned} \tag{4.1}$$

hold. Now, let  $\Psi'(x)$  be associated with any new basis

$$\Psi'(x) = S \Psi(x), \quad \Gamma'^a = S \Gamma^a S^{-1}, \tag{4.2a}$$

then

$$C' (\Gamma'^a)^* (C')^{-1} = +\Gamma'^a \tag{4.2b}$$

and further

$$S^{-1} C' S^* (\Gamma^a)^* S^{-1} (C')^{-1} S = +\Gamma^a \implies S^{-1} C' S^* = C.$$

Thus, a rule for transforming  $C$ -matrix is

$$C' = S C (S^*)^{-1}. \quad (4.2c)$$

Now, let us follow up in some detail how, having started from  $\Psi'''$ -basis (when  $C''' = I$ , see (2.6)), one can reconstruct  $C$ -operation in  $\Psi''$ -representation:

$$\Psi'' = \begin{pmatrix} d_0 & 0 & 0 & 0 \\ 0 & d_0 & 0 & 0 \\ 0 & 0 & I & 0 \\ 0 & 0 & 0 & I \end{pmatrix} \Psi''' \quad \Rightarrow \quad C'' = I. \quad (4.3)$$

Next step is

$$\Psi' = \begin{pmatrix} e^{+it} & 0 & 0 & 0 \\ 0 & e^{+it} & 0 & 0 \\ 0 & 0 & I & 0 \\ 0 & 0 & 0 & I \end{pmatrix} \Psi'' \quad \Rightarrow \quad C' = \begin{pmatrix} (d/d^*) & 0 & 0 & 0 \\ 0 & (d/d^*) & 0 & 0 \\ 0 & 0 & I & 0 \\ 0 & 0 & 0 & I \end{pmatrix}. \quad (4.4)$$

And last step is

$$\Psi = \begin{pmatrix} 1 & \lambda_4^*/D^* & \lambda_2/D^* & 0 \\ 0 & \lambda_3^*/D^* & -\lambda_1/D^* & 0 \\ 0 & 0 & 0 & I \end{pmatrix} \Psi' \quad \Rightarrow \quad C = \begin{pmatrix} d/d^* & 0 & 0 & 0 \\ 0 & [\lambda_1^* \lambda_4^* d + \lambda_2 \lambda_3 d^*]/d^* D^* & [\lambda_2^* \lambda_4^* d - \lambda_2 \lambda_4 d^*]/d^* D^* & 0 \\ 0 & [\lambda_1^* \lambda_3^* d - \lambda_1 \lambda_3 d^*]/d^* D^* & [\lambda_2^* \lambda_3^* d + \lambda_1 \lambda_4 d^*]/d^* D^* & 0 \\ 0 & 0 & 0 & I \end{pmatrix}. \quad (4.5)$$

It should be noted that the central two-by-two block is quite symmetric one:

$$\begin{aligned} (\lambda_2^* \lambda_4^* d - \lambda_2 \lambda_4 d^*) &= (\lambda_1 \lambda_2^* \lambda_3 \lambda_4^* - \lambda_1^* \lambda_2 \lambda_3^* \lambda_4), \\ (\lambda_1^* \lambda_3^* d - \lambda_1 \lambda_3 d^*) &= -(\lambda_1 \lambda_2^* \lambda_3 \lambda_4^* - \lambda_1^* \lambda_2 \lambda_3^* \lambda_4). \end{aligned} \quad (4.6)$$

Now let us apply else one method for establishing the same  $C$ -matrix: via a direct analysis of the defining relation

$$C(\Gamma^a)^* C^{-1} = +\Gamma^a \quad \Rightarrow \quad C(\Gamma^a)^* = \Gamma^a C. \quad (4.7a)$$

Taking  $C$  in the form (see (4.3))

$$C = \begin{pmatrix} P_1 & 0 & 0 & 0 \\ 0 & P_2 & P_3 & 0 \\ 0 & P_4 & P_5 & 0 \\ 0 & 0 & 0 & P_6 \end{pmatrix},$$

where  $P_1$  is a number,  $P_2, P_3, P_4, P_5$  and  $P_6$  are proportional to a four-by-four and a six-by-six identity matrices respectively, we will have

$$\begin{aligned} \lambda_1^* P_1 &= \lambda_1 P_2^* + \lambda_2 P_4^* & \lambda_2^* P_1 &= \lambda_1 P_3^* + \lambda_2 P_5^* \\ \lambda_3^* P_6 &= \lambda_3^* P_2^* - \lambda_4^* P_4^* & -\lambda_4 P_6 &= \lambda_3^* P_3^* - \lambda_4^* P_5^* \end{aligned}, \quad (4.8a)$$

$$\begin{aligned} -\lambda_1^* P_1^* &= -\lambda_1 P_2 + \lambda_2 P_3 & \lambda_2^* P_1^* &= -\lambda_1 P_4 + \lambda_2 P_5 \\ \lambda_3 P_6^* &= \lambda_3^* P_2 + \lambda_4^* P_3 & \lambda_4 P_6^* &= \lambda_3^* P_4 + \lambda_4^* P_5. \end{aligned} \quad (4.8b)$$

One can simplify the task by taking yet known forms for  $P_1$  and  $P_6$ :  $P_1 = d/d^*$ ,  $P_6 = 1$ . Then eqs. (4.8b) lead to

$$\begin{aligned} \lambda_1^* \frac{d}{d^*} &= \lambda_1 P_2^* + \lambda_2 P_4^* & \lambda_2^* \frac{d}{d^*} &= \lambda_1 P_3^* + \lambda_2 P_5^* \\ \lambda_3 &= \lambda_3^* P_2^* - \lambda_4^* P_4^* & -\lambda_4 &= \lambda_3^* P_3^* - \lambda_4^* P_5^* \end{aligned}, \quad (4.9)$$

$$\begin{aligned} -\lambda_1^* \frac{d^*}{d} &= -\lambda_1 P_2 + \lambda_2 P_3 & \lambda_2^* \frac{d^*}{d} &= -\lambda_1 P_4 + \lambda_2 P_5 \\ \lambda_3 &= \lambda_3^* P_2 + \lambda_4^* P_3 & \lambda_4 &= \lambda_3^* P_4 + \lambda_4^* P_5 \end{aligned} \quad (4.10)$$

and their solution follows quite straightforwardly

$$P_2 = \frac{d^* \lambda_2 \lambda_3 + d \lambda_1^* \lambda_4^*}{(\lambda_1 \lambda_4^* + \lambda_2 \lambda_3^*) d^*}, \quad P_3 = \frac{\lambda_1 \lambda_2^* \lambda_3 \lambda_4^* - \lambda_1^* \lambda_2 \lambda_3^* \lambda_4}{(\lambda_1 \lambda_4^* + \lambda_2 \lambda_3^*) d^*}, \quad (4.11a)$$

$$P_5 = \frac{d \lambda_2^* \lambda_3^* + d^* \lambda_1^* \lambda_4}{(\lambda_1 \lambda_4^* + \lambda_2 \lambda_3^*) d^*}, \quad P_4 = \frac{\lambda_1^* \lambda_2 \lambda_3^* \lambda_4 - \lambda_1 \lambda_2^* \lambda_3 \lambda_4^*}{(\lambda_1 \lambda_4^* + \lambda_2 \lambda_3^*) d^*}, \quad (4.11b)$$

what coincides with (4.5).

As known, a knowledge of the  $C$ -matrix provides us with a possibility to reconstruct an anti-particle's wave functions in terms of those of particles. Also, it enables us to isolate chargeless states  $\Psi^\pm$  from charged ones  $\Psi$ :

$$\Psi^c = C^* \Psi = C \Psi^*, \quad \Psi^\pm = \frac{1}{2}(\Psi \pm \Psi^c), \quad C^* \Psi^\pm = (\pm 1) \Psi^\pm. \quad (4.12a)$$

Let us remember that the  $U(1)$ -gauge invariant derivative  $(\partial_a - ieA_a)$  necessarily mixes up together  $\Psi^+$  and  $\Psi^-$  constituents. Really, combining two equations

$$[\Gamma^a(\partial_a - ieA_a) - m] \Psi = 0, \quad [\Gamma^a(\partial_a + ieA_a) - m] \Psi^c = 0,$$

one can readily produce

$$\Gamma^a(\partial_a \Psi^\pm - igA_a \Psi^\mp) - m \Psi^\pm = 0. \quad (4.12b)$$

## 5. Equations in a Riemannian space-time, tensor-based approach

Section 5 deals with equations for the spin 1 particle adjusted to the general relativity requirements. We are going to start with general covariant tensor formalism; later, in Sec. 7, an alternative approach, based on the tetrad<sup>4</sup> method by Tetrad-Weyl-Fock-Ivanenko [46-80], will be developed in full detail. Such a study could disclose how the additional  $\sigma$ -characteristic can exhibit itself at the background of non-Euclidean geometry. Besides, the generally covariant formalism will enable us to check correctness of the above determined massless limit of the theory: reminding that massless particle's covariant equations are assumed to be conformally invariant.

On taking a simple formal change of Cartesian derivatives and tensors into covariant ones

$$\partial_b \Rightarrow \nabla_\beta, \quad C_b(x) \Rightarrow C_\beta(x), \quad \Phi_b(x) \Rightarrow \Phi_\beta(x), \quad \Phi_{ab}(x) \Rightarrow \Phi_{\alpha\beta}(x)$$

eqs. (2.6) take on the form

$$\begin{aligned} \nabla^\alpha C_\alpha - mC &= 0, \quad \nabla^\alpha \Phi_{\beta\alpha} - mC_\beta = 0, \\ \sigma \nabla_\beta C + \nabla^\alpha \Phi_{\beta\alpha} - m\Phi_\beta &= 0, \\ \nabla_\alpha \Phi_\beta - \nabla_\beta \Phi_\alpha - m\Phi_{\alpha\beta} &= 0. \end{aligned} \quad (5.1)$$

Obviously, in presence of external electromagnetic fields, with the notation  $\nabla_\alpha \Rightarrow D_\alpha = \nabla_\alpha - igA_\alpha(x)$ , instead of (5.1) we will have

$$\begin{aligned} D^\alpha C_\alpha - mC &= 0, \quad D^\alpha \Phi_{\beta\alpha} - mC_\beta = 0, \\ \sigma D_\beta C + D^\alpha \Phi_{\beta\alpha} - m\Phi_\beta &= 0, \\ D_\alpha \Phi_\beta - D_\beta \Phi_\alpha - m\Phi_{\alpha\beta} &= 0. \end{aligned} \quad (5.2)$$

Let us exclude the field  $C$  from third equation in (5.2). To this end, one can produce

$$D^\alpha C_\alpha = \frac{1}{m} D^\alpha D^\beta \Phi_{\alpha\beta}, \quad D^\alpha D^\beta \Phi_{\alpha\beta} = \nabla^\alpha \nabla^\beta \Phi_{\alpha\beta} - ig \frac{1}{2} F^{\alpha\beta} \Phi_{\alpha\beta},$$

where  $F_{\alpha\beta} = (\nabla_\alpha A_\beta - \nabla_\beta A_\alpha)$ , and further

$$\nabla^\alpha \nabla^\beta \Phi_{\alpha\beta} = \frac{1}{2} (\nabla^\alpha \nabla^\beta - \nabla^\beta \nabla^\alpha) \Phi_{\alpha\beta} =$$

---

<sup>4</sup>Below, both terms, tetrad and vierbein, will be used interchangeably.

$$= \frac{1}{2} [ \Phi_{\nu\alpha}(x) R^{\nu\beta\alpha}(x) + \Phi_{\beta\nu}(x) R^{\nu\beta\alpha}(x) ] = \frac{1}{2} [ \Phi_{\nu\alpha}(x) R^{\nu\alpha}(x) + \Phi_{\nu\beta}(x) R^{\nu\beta}(x) ] .$$

Now, remembering symmetry of the Ricci tensor  $R_{\alpha\beta}(x)$ , we arrive at

$$D^\alpha C_\alpha = +i \frac{g}{m} \frac{1}{2} F^{\alpha\beta} \Phi_{\alpha\beta} .$$

Since, first equation in (5.2) yields

$$C = -i \frac{g}{m^2} \frac{1}{2} F^{\alpha\beta} \Phi_{\alpha\beta} . \quad (5.3)$$

Therefore, the main equations in (5.2) read as

$$\begin{aligned} D^\alpha C_\alpha - mC &= 0 , & D^\alpha \Phi_{\beta\alpha} - mC_\beta &= 0 , \\ -i\sigma \frac{g}{m^2} D_\rho \left( \frac{1}{2} F^{\alpha\beta} \Phi_{\alpha\beta} \right) + D^\alpha \Phi_{\rho\alpha} - m\Phi_\rho &= 0 , \\ D_\alpha \Phi_\beta - D_\beta \Phi_\alpha - m\Phi_{\alpha\beta} &= 0 . \end{aligned} \quad (5.4)$$

In absence external electromagnetic fields, the term proportional to  $\sigma$  vanishes. This means that  $\sigma$ -characteristics manifests itself physically only in electromagnetic fields but not in gravitational ones.

Now, let us consider a generally covariant massless equations. They are to be as follows

$$\begin{aligned} \nabla^\alpha C_\alpha - C &= 0 , & \nabla^\alpha \Phi_{\beta\alpha} - C_\beta &= 0 , \\ \sigma \nabla_\beta C + \nabla^\alpha \Phi_{\beta\alpha} &= 0 , & \nabla_\alpha \Phi_\beta - \nabla_\beta \Phi_\alpha - \Phi_{\alpha\beta} &= 0 . \end{aligned} \quad (5.5)$$

As in the flat space-time, here we have

$$\begin{aligned} C &= 0 , & C_\beta &= 0 , \\ \nabla^\alpha \Phi_{\beta\alpha} &= 0 , & \nabla_\alpha \Phi_\beta - \nabla_\beta \Phi_\alpha - \Phi_{\alpha\beta} &= 0 . \end{aligned} \quad (5.6)$$

Thus, eqs. (5.6) happen to be absolutely equivalent to ordinary generally covariant Proca massless equations.

Repeating all arguments after formula (1.18) from Sec. 1, for a generally covariant massless particle in an external vector field, we will get

$$\begin{aligned} C_\beta &= D^\alpha \Phi_{\alpha\beta} , & C &= -ig \frac{1}{2} F^{\alpha\beta} \Phi_{\alpha\beta} , \\ -i\sigma g D_\rho \left( \frac{1}{2} F^{\alpha\beta} \Phi_{\alpha\beta} \right) + D^\alpha \Phi_{\rho\alpha} &= 0 , \\ D_\alpha \Phi_\beta - D_\beta \Phi_\alpha - \Phi_{\alpha\beta} &= 0 . \end{aligned} \quad (5.7)$$

## 6. Covariant tetrad-based formalism

We start with the matrix equation in the Minkowski space-time (2.12)

$$(\Gamma^a \partial_a - m) \Psi(x) = 0 , \quad (6.1a)$$

$$\Psi = (C, C_a, \Phi_a, \Phi_{ab}), \quad \Gamma^a = \begin{pmatrix} 0 & G^a & 0 & 0 \\ 0 & 0 & 0 & K^a \\ \sigma \Delta^a & 0 & 0 & K^a \\ 0 & 0 & \Lambda^a & 0 \end{pmatrix} , \quad (6.1b)$$

$$\begin{aligned} (K^a)_l{}^{mn} &= (-g^{am} \delta_l^n + g^{an} \delta_l^m) , \\ (\Lambda^a)_{bl}{}^k &= \delta_{bl}^{ak} , \quad (G^a)_{(0)}^k = g^{ab} , \quad (\Delta^a)_l{}^{(0)} = \delta_l^a . \end{aligned} \quad (6.1c)$$

In accordance with the known procedure by Tetrode-Weyl-Fock-Ivanenko, eq. (6.1) is extended to a curved space-time with a metric tensor  $g_{\alpha\beta}(x)$  and an accompanying tetrad  $e_{(a)}^\alpha(x)$  as follows

$$[\Gamma^\alpha(x) (\partial_\alpha + B_\alpha(x)) - m] \Psi(x) = 0 , \quad (6.2)$$



where

$$\Gamma^\alpha(x) = \Gamma^a e_{(a)}^\alpha(x) , \quad B_\alpha(x) = \frac{1}{2} J^{ab} e_{(a)}^\beta \nabla_\alpha (e_{(b)\beta}) .$$

Here  $J^{ab}$  designate generators of the Lorentz group representation (see below formulas (6.6)) associated with the set  $(C, C_k, \Phi_k, \Phi_{kl})$ .

The equation (6.2) involves the object  $e_{(a)}^\alpha(x)$ . Therefore, similar to the case of much more familiar spin 1/2 particle, there must exist possibility that any such equations associated respectively with  $e_{(a)}^\alpha(x)$  and  $e'_{(a)}^\alpha(x)$  were translated into each other; otherwise eq. (6.2) is incorrect.

Let us show that if two tetrads are connected by a Lorentz operator  $L$ :

$$e_{(a)}^\alpha(x) \implies e'_{(b)}^\alpha(x) = L_a^b(x) e_{(b)}^\alpha(x) , \quad (6.3a)$$

the respective two equations

$$\begin{aligned} [\Gamma^\alpha(x)(\partial_\alpha + B_\alpha(x)) - m] \Psi(x) &= 0 , \\ [\Gamma'^\alpha(x)(\partial_\alpha + B'_\alpha(x)) - m] \Psi'(x) &= 0 \end{aligned} \quad (6.3b)$$

can be set into each other by means of a Lorentz gauge transformation

$$\Psi'(x) = S(x) \Psi(x) , \quad \begin{pmatrix} C'(x) \\ C'_k(x) \\ \Phi'_k(x) \\ \Phi'_{kl}(x) \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & L_k^l & 0 & 0 \\ 0 & 0 & L_k^l & 0 \\ 0 & 0 & 0 & L_k^m L_l^n \end{pmatrix} \begin{pmatrix} C(x) \\ C_l(x) \\ \Phi_l(x) \\ \Phi_{mn}(x) \end{pmatrix} . \quad (6.3c)$$

It is useful time to dwell upon some peculiarities of the formalism. The form of the  $15 \times 15$ -gauge transformation (6.3c) by no means takes into consideration antisymmetry of the tensor  $\Phi_{kl}$ . If such a symmetry is allowed for, then the Lorentz operator can be rewritten in a modified form

$$\Phi'_{kl}(x) = L_k^m L_l^n \Phi_{mn} = \frac{1}{2} (L_k^m L_l^n - L_l^m L_k^n) \Phi_{mn} . \quad (6.4a)$$

This formula (6.4a) has some advantage in the context of the matrix formalism, in view of presence of the tensor  $\Phi_{mn}$  just as a 6-dimension object  $(\Phi_{01}, \Phi_{02}, \Phi_{03}, \Phi_{23}, \Phi_{31}, \Phi_{12})$ . At this the 6-vector transforms by means of the operator

$$S_{6 \otimes 6} = S_{kl}{}^{mn} = (L_k^m L_l^n - L_l^m L_k^n) , \quad S_{6 \otimes 6} = L_{[} \otimes L_{]} . \quad (6.4b)$$

The property noted plays a role and for corresponding generators. Let  $V^{ab}$  and  $(V \otimes V)^{ab}$  designate respectively vector and tensor generators:

$$\Phi'_s = (\delta_s^p + \frac{1}{2} \delta \omega_{ab} (V^{ab})_s{}^p) \Phi_p , \quad (V^{ab})_s{}^p = -g^{ap} \delta_s^b + g^{bp} \delta_s^a , \quad (6.5a)$$

$$\begin{aligned} \Phi'_{mn} &= ((\delta_m^s + \frac{1}{2} \delta \omega_{ab} (V^{ab})_m{}^s) (\delta_n^p + \frac{1}{2} \delta \omega_{ab} (V^{ab})_n{}^p) \Phi_{sp} = \\ &= \delta_m^s \delta_n^p + \frac{1}{2} \delta \omega_{ab} [(V \otimes V)^{ab}]_{mn}{}^{sp} \Phi_{sp} , \end{aligned}$$

where

$$[(V \otimes V)^{ab}]_{mn}{}^{sp} = [(-g^{as} \delta_m^b + g^{bs} \delta_m^a) \delta_n^p + \delta_m^s (-g^{ap} \delta_n^b + g^{bp} \delta_n^a)] , \quad (6.5b)$$

or

$$(V \otimes V)^{ab} = V^{ab} \otimes I + I \otimes V^{ab} .$$

Here the tensor generator does not take into consideration any symmetry property of  $\Phi_{sp}$ , otherwise we will have

$$\begin{aligned} [V^{ab}]_{mn}{}^{sp} \Phi_{sp} &= \frac{1}{2} \{ [(V \otimes V)^{ab}]_{mn}{}^{sp} - [(V \otimes V)^{ab}]_{mn}{}^{ps} \} \Phi_{sp} = \\ &= \frac{1}{2} \{ (-\delta_{mn}^{as} g^{bp} + \delta_{mn}^{ap} g^{bs}) - (-\delta_{mn}^{bs} g^{ap} + \delta_{mn}^{bp} g^{as}) \} \Phi_{sp} . \end{aligned} \quad (6.5)$$

Thus, there exist two usages of the generators  $J^{abn}$  and gauge transformation  $S$ :

$$J^{ab} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & V^{ab} & 0 & 0 \\ 0 & 0 & V^{ab} & 0 \\ 0 & 0 & 0 & (V \otimes V)^{ab} \end{pmatrix}, \quad S = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & L & 0 & 0 \\ 0 & 0 & L & 0 \\ 0 & 0 & 0 & L \otimes L \end{pmatrix}, \quad (6.6a)$$

or

$$J_{[]}^{ab} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & V^{ab} & 0 & 0 \\ 0 & 0 & V^{ab} & 0 \\ 0 & 0 & 0 & (V_{[]} )^{ab} \end{pmatrix}, \quad S_{[]} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & L & 0 & 0 \\ 0 & 0 & L & 0 \\ 0 & 0 & 0 & L_{[]} \otimes L_{[]} \end{pmatrix} \quad (6.6b)$$

Now, starting from an equation for  $\Psi$ -function, let us obtain an equation for  $\Psi'$ -function. We have

$$[ S \Gamma^\alpha S^{-1} (\partial_\alpha + S B_\alpha S^{-1} + S \partial_\alpha S^{-1}) - M ] \Psi' = 0,$$

and two relationships are to be proved

$$S \Gamma^\alpha S^{-1} = \Gamma'^\alpha, \quad (6.7a)$$

$$S B_\alpha S^{-1} + S \partial_\alpha S^{-1} = B'_\alpha. \quad (6.7b)$$

The first (6.7a) can be rewritten as

$$S \Gamma^a e_{(a)}^\alpha S^{-1} = \Gamma^b e_{(b)}'^\alpha \implies S \Gamma^a S^{-1} = \Gamma^b L_b^a. \quad (6.8a)$$

The last is a well-known condition insuring the relativistic invariance property of the 15-component wave equation being considered in the Minkowski space. In blocks it reads

$$G^a L^{-1} = G^b L_b^a, \quad L \Delta^a = \Delta^b L_b^a, \\ L K^a (L^{-1} \otimes L^{-1}) = K^b L_b^a, \quad (L \otimes L) \Lambda^a L^{-1} = \Lambda^b L_b^a. \quad (6.8b)$$

One can check these relations by a direct calculation with the use of the explicit form of all the blocks  $G^a, \Delta^a, K^a, \Lambda^a$ . The first one gives

$$(G^a)_{(0)}^k (L^{-1})_k^c = (G^b)_{(0)}^c L_b^a \implies (L^{-1})^{ac} = L^{ca},$$

which coincides with the known pseudo orthogonality property of the Lorentz matrix. Second one leads to an identity

$$L_k^l (\Delta^a)_l^{(0)} = (\Delta^b)_k^{(0)} L_b^a \implies L_k^a = L_k^a;$$

Third yields

$$L_d^l (K^a)_l^{mn} [(L^{-1})_m^s (L^{-1})_n^p] = (K^b)_d^{sp} L_b^a,$$

what is equivalent to

$$\delta_d^p (L^{-1})^{as} - \delta_d^s (L^{-1})^{ap} = \delta_d^p L^{sa} - \delta_d^s L^{pa}.$$

And finally forth reads as

$$L_m^c L_n^d (\Lambda^a)_{cd}^k (L^{-1})_k^l = \Lambda^b L_b^a$$

that is an identity

$$\delta_n^l L_m^a - \delta_m^l L_n^a = \delta_n^l L_m^a - \delta_m^l L_n^a.$$

Now let us turn to eq. (6.7b) and consider in some detail the term

$$S B_\alpha S^{-1} = \frac{1}{2} J^{ab} S^{-1} e_{(a)}^\beta (\nabla_\alpha e_{(b)\beta}). \quad (6.9)$$

Expressing the vierbein  $e_{(b)\beta}$  in terms of the primed  $e'_{(b)\beta}$ , we get

$$S B_\alpha S^{-1} = \frac{1}{2} (S J^{ab} S^{-1}) (L^{-1})_a^k e_{(k)}'^\beta [\nabla_\alpha (L^{-1})_b^l e_{(l)\beta}]$$

and further

$$S B_\alpha S^{-1} = \frac{1}{2} (S J^{ab} S^{-1}) \times$$

$$\begin{aligned}
& \times \left[ (L^{-1})_a{}^k e'_{(k)}{}^\beta \left[ (\partial_\alpha (L^{-1})_b{}^l) e'_{(l)\beta} + (L^{-1})_b{}^l \nabla_\alpha e'_{(l)\beta} \right] = \right. \\
& = \frac{1}{2} (S J^{ab} S^{-1}) \left[ (L^{-1})_a{}^k (\partial_\alpha (L^{-1})_b{}^l) e'_{(k)}{}^\beta e'_{(l)\beta} + (L^{-1})_a{}^k (L^{-1})_b{}^l e'_{(k)}{}^\beta \nabla_\alpha e'_{(l)\beta} \right].
\end{aligned}$$

So, one arrives at

$$\begin{aligned}
S B_\alpha S^{-1} &= \frac{1}{2} (S J^{ab} S^{-1}) \times \\
&\times \left[ (L^{-1})_a{}^k (\partial_\alpha (L^{-1})_b{}^l) g_{kl} + (L^{-1})_a{}^k (L^{-1})_b{}^l e'_{(k)}{}^\beta \nabla_\alpha e'_{(l)\beta} \right].
\end{aligned} \tag{6.10}$$

Taking into account the identity (its proof will be given below)

$$S J^{ab} S^{-1} = J^{mn} L_m{}^a L_n{}^b, \tag{6.11}$$

from (6.10) one readily obtains

$$\begin{aligned}
S B_\alpha S^{-1} &= \frac{1}{2} J^{mn} L_n{}^b \partial_\alpha (L^{-1})_{bm} + \frac{1}{2} J^{kl} e'_{(k)}{}^\beta \nabla_\alpha e'_{(l)\beta} = \\
&= \frac{1}{2} J^{mn} L_n{}^b \partial_\alpha (L^{-1})_{bm} + B'_\alpha.
\end{aligned} \tag{6.12}$$

Now, on taking into consideration (6.12), from (6.7b) it follows

$$S \partial_\alpha S^{-1} + \frac{1}{2} J^{mn} L_n{}^b \partial_\alpha (L^{-1})_{bm} = 0$$

which can be rewritten as

$$S \partial_\alpha S^{-1} = \frac{1}{2} J^{mn} L_{mb} (\partial_\alpha L_n{}^b). \tag{6.13}$$

With the use of all blocks involved, the previous relation will read as

$$L \partial_\beta L^{-1} = \frac{1}{2} V^{ab} L_{ad} \partial_\beta L_b{}^d, \tag{6.14}$$

$$(L^{-1} \otimes L^{-1}) \partial_\beta (L^{-1} \otimes L^{-1}) = \frac{1}{2} (V \otimes V)^{ab} L_{ad} \partial_\beta L_b{}^d. \tag{6.14b}$$

It suffices to prove only (6.14a), because eq. (6.14b) is a straightforward result from that on using the formula  $(V \otimes V)^{ab} = (V^{ab} \otimes I + I \otimes V^{ab})$ . Eq. (6.14a) yields

$$L_s{}^k \partial_\beta (L^{-1})_k{}^p = \frac{1}{2} (g^{ap} \delta_s^b - g^{bp} \delta_s^a) L_{ad} \partial_\beta L_b{}^d,$$

or

$$L_{sd} \partial_\beta (L^{-1})^{dp} = \frac{1}{2} [L^{pd} \partial_\beta L_{sd} - L_{sd} \partial_\beta L^{pd}].$$

Here a supplementary relation is needed, which may be resulted in by differentiating the known pseudo-orthogonality relation for  $L$ :

$$\partial_\beta [L_{sd} L^{pd}] = (\partial_\beta L_{sd}) L^{pd} + L_{sd} \partial_\beta L^{pd} = 0 \tag{6.15}$$

With the use of eq. (6.15) the previous one reads as identity. So, eq. (6.13) has been proved. .

Now, we are to return to eq. (6.11) used above. It is equivalent to

$$L V^{ab} L^{-1} = V^{mn} L_m{}^a L_n{}^b, \tag{6.16a}$$

$$(L \otimes L)(V \otimes V)(L^{-1} \otimes L^{-1}) = (V \otimes V)^{mn} L_m{}^a L_n{}^b. \tag{6.16b}$$

Obviously, the second is a simple consequence of the first. In turn, the first one (6.16a) happens to be an identity:

$$\begin{aligned}
L(V^{ab})_s{}^p L^{-1} &= (V^{mn})_s{}^p L_m{}^a L_n{}^b \implies \\
-L_k{}^b L^{sa} + L_k{}^a + L_k{}^a L^{sb} &= -L_k{}^b L^{sa} + L_k{}^a + L_k{}^a L^{sb}.
\end{aligned}$$

Thus, any two equations (6.36) associated with two different tetrads are transforms of each other. This means that the gauge invariance principle with respect to local Lorentz group, in a manner like more familiar case of spin 1/2 particle, holds.

Else one fact should be mentioned. The wave equation under consideration will look as correct one in a generally relativistic context, if the 15-component wave function  $\Psi(x)$  is a scalar with respect to general coordinate transformations:

$$x'^\alpha = f^\alpha(x^\alpha) \implies \Psi'(x') = \Psi(x) . \quad (6.17)$$

Now let us show — how tensor equations discussed in Sec. 5 are resulted in from the matrix one:

$$[ \Gamma^\rho(x)(\partial + B_\rho(x)) - m ] \Psi(x) = 0 . \quad (6.18)$$

To this end, as a first step, let us take into consideration the block structure of all objects involved:

$$\begin{aligned} \Gamma^\rho(x) = e_{(l)}^\rho \Gamma^l &= \begin{pmatrix} 0 & G^\rho & 0 & 0 \\ 0 & 0 & 0 & K^\rho \\ \sigma \Delta^\rho & 0 & 0 & K^\rho \\ 0 & 0 & \Lambda^\rho & 0 \end{pmatrix} , \quad B_\rho(x) = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & V_\rho & 0 & 0 \\ 0 & 0 & V_\rho & 0 \\ 0 & 0 & 0 & (V \otimes V)_\rho \end{pmatrix} , \\ \Gamma^\rho(\partial_\rho + B_\rho) &= \begin{pmatrix} 0 & G^\rho[\partial_\rho + V_\rho] & 0 & 0 \\ 0 & 0 & 0 & K^\rho[\partial_\rho + (V \otimes V)_\rho] \\ \sigma \Delta^\rho \partial_\rho & 0 & 0 & K^\rho[\partial_\rho + (V \otimes V)_\rho] \\ 0 & 0 & \Lambda^\rho[\partial_\rho + V_\rho] & 0 \end{pmatrix} . \end{aligned} \quad (6.19)$$

Below it will convenient to employ the notation

$$(V_\rho)_l{}^k(x) = \frac{1}{2}(V^{ab})_l{}^k e_{(a)}^\beta \nabla_\rho e_{(b)\beta} , \quad (6.20)$$

$$[(V \otimes V)_{\rho}]_{ps}{}^{mn}(x) = \frac{1}{2}[(V \otimes V)^{ab}]_{ps}{}^{mn} e_{(a)}^\beta \nabla_\rho e_{(b)\beta} .$$

Then eq.(6.18) takes on the block form

$$\begin{aligned} [ G^\rho (\partial_\rho + V_\rho) ]_{(0)}^k C_k &= mC , \\ \{ K^\rho [\partial_\rho + (V \otimes V)_\rho] \}_l{}^{mn} \Phi_{mn} &= mC_l , \\ \sigma (\Delta^\rho)_{(l)}^{(0)} \partial_\rho C + \{ K^\rho [\partial_\rho + (V \otimes V)_\rho] \}_l{}^{mn} \Phi_{mn} &= m\Phi_l , \\ [ \Lambda^\rho (\partial_\rho + V_\rho) ]_{mn}^k \Phi_k &= m\Phi_{mn} . \end{aligned} \quad (6.21)$$

With the use of explicit form of the all block-matrices in (6.21) one can readily arrive at

$$\begin{aligned} e^{(k)\alpha} \partial_\alpha C_k + \gamma_l{}^{kl} C_k &= mC , \\ e^{(k)\alpha} \partial_\alpha \Phi_{lk} + \gamma_l{}^{mn} \Phi_{mn} + \gamma_d{}^{kd} \Phi_{lk} &= mC_l , \\ \sigma e_{(l)}^\alpha \partial_\alpha C + e^{(k)\alpha} \partial_\alpha \Phi_{lk} + \gamma_l{}^{mn} \Phi_{mn} + \gamma_d{}^{kd} \Phi_{lk} &= m\Phi_l , \\ e_{(m)}^\alpha \partial_\alpha \Phi_n - e_{(n)}^\alpha \partial_\alpha \Phi_m + (\gamma_{mn}^k - \gamma_{nm}^k) \Phi_k &= m\Phi_{mn} . \end{aligned} \quad (6.22)$$

Here  $\gamma_{abc}$  designates the Ricci rotation coefficients

$$\gamma_{abc}(x) = -e_{(a)\rho;\sigma} e_{(b)}^\rho e_{(c)}^\sigma .$$

In turn, as could be checked, eqs. (6.22) represent the following generally covariant tensor equations (just those introduced in Sec. 5)

$$\begin{aligned} \nabla^\alpha C_\alpha - mC &= 0 , \quad \nabla^\alpha \Phi_{\beta\alpha} - mC_\beta = 0 , \\ \sigma \nabla_\beta C + \nabla^\alpha \Phi_{\beta\alpha} - m\Phi_\beta &= 0 , \quad \nabla_\alpha \Phi_\beta - \nabla_\beta \Phi_\alpha - m\Phi_{\alpha\beta} = 0 , \end{aligned} \quad (6.23)$$

in the vierbein form. Connection between generally covariant and tetrad representatives of the wave function is determined quite conventionally

$$C_\alpha = e_\alpha^{(l)} C_l , \quad \Phi_\alpha = e_\alpha^{(l)} \Phi_l , \quad \Phi_{\alpha\beta} = e_\alpha^{(m)} e_\beta^{(n)} \Phi_{mn} . \quad (6.24)$$

## 7. Bilinear invariants in a curved space-time

This Section deals with a generally covariant equation for a  $\bar{\Psi}$ -function conjugate to the  $\Psi$  and methods for constructing some bilinear invariants in terms of  $\bar{\Psi}$  and  $\Psi$ . A knowing of this matter will enable us to produce automatically expressions for a Lagrangian and a conserved current (about the energy-momentum tensor see Sec. 9).

The basic requirement of an invariance matrix  $\eta$  is that a form  $\Psi^+ \eta \Psi$  be invariant under Lorentz group transformations

$$\Psi^+ \eta \Psi = inv, \quad \Psi' = S \Psi,$$

which necessitates

$$\eta = S^+ \eta S \quad \implies \quad \eta S^{-1} = S^+ \eta. \quad (7.1a)$$

Being taken for an infinitesimal  $S$ , eq. (7.1a) reads

$$-\eta J^{ab} = (J^{ab})^+ \eta; \quad (7.1b)$$

it is what we need. Now, starting from an equation for  $\Psi$

$$[\Gamma^\alpha (\partial_\alpha + B_\alpha) - m] \Psi = 0, \quad (7.2)$$

we are going to reconstruct that for  $\bar{\Psi} = \Psi^+ \eta$ . From eq. (7.2) it follows

$$\Psi^+ \eta \eta^{-1} [(\overleftarrow{\partial}_\alpha + B_\alpha^+) \eta \eta^{-1} (\Gamma^\alpha)^+ - m] \eta = 0;$$

which yields

$$\eta^{-1} [\Gamma^\alpha(x)]^+ \eta = -\Gamma^\alpha(x), \quad \eta^{-1} [B_\alpha(x)]^+ \eta = -B_\alpha(x), \quad (7.3)$$

and further (compare with (7.2))

$$\bar{\Psi} [(\overleftarrow{\partial}_\alpha - B_\alpha) \Gamma^\alpha + M] = 0. \quad (7.4)$$

Take notice on the sign 'minus' at the spinor connection  $B_\alpha$  in (7.4). Now, multiplying eq. (7.2) from the left by  $\bar{\Psi}$ , and eq. (7.4) — from the right by  $\Psi$ , and adding results together, we get to

$$\begin{aligned} & \bar{\Psi} \overleftarrow{\partial}_\alpha \Gamma^\alpha \Psi + \bar{\Psi} \Gamma^\alpha \overrightarrow{\partial}_\alpha \Psi + \\ & + \bar{\Psi} (\Gamma^\alpha B_\alpha - B_\alpha \Gamma^\alpha) \Psi = 0. \end{aligned} \quad (7.5)$$

To proceed with an analysis of eq. (7.5), we need one auxiliary relation. To this end, let us turn again to the used above formula  $S \Gamma^a S^{-1} = \Gamma^b L_b^a$  and take it for an infinitesimal Lorentz transformation:

$$(1 + \frac{1}{2} \delta \omega_{mn} J^{mn}) \Gamma^a (1 - \frac{1}{2} \delta \omega_{kl} J^{kl}) = \Gamma^b [\delta_b^a + \frac{1}{2} \delta \omega_{kl} (V^{kl})_b^a],$$

from where it follows

$$J^{kl} \Gamma^a - \Gamma^a J^{kl} = \Gamma^b (V^{kl})_b^a.$$

Taking into account the generators  $V^{kl}$  explicitly we will have the commutation relation

$$J^{kl} \Gamma^a - \Gamma^a J^{kl} = \Gamma^k g^{lk} - \Gamma^l g^{ka}. \quad (7.6)$$

Now, multiplying eq. (7.6) by an expression  $e_{(a)}^\rho \frac{1}{2} e_{(k)}^\beta \nabla_\sigma e_{(l)\beta}$  we arrive at

$$\Gamma^\rho B_\sigma - B_\sigma \Gamma^\rho = \nabla_\sigma \Gamma^\rho = \Gamma^\rho_{;\sigma}. \quad (7.7)$$

Including (7.7) in (7.5), we come to

$$\bar{\Psi} \overleftarrow{\partial}_\alpha \Gamma^\alpha \Psi + \bar{\Psi} \Gamma^\alpha \overrightarrow{\partial}_\alpha \Psi + \bar{\Psi} (\nabla_\alpha \Gamma^\alpha) \Psi = 0,$$

which may be rewritten as a generally covariant conserved current law:

$$\nabla_\alpha J^\alpha = 0, \quad J^\alpha = \bar{\Psi} \Gamma^\alpha \Psi. \quad (7.8a)$$

The conserved current  $J^\alpha(x)$  may be expressed in terms of generally covariant tensor components according to (see (3.20); a numerical factor is omitted)

$$J^\alpha = +\sigma ( C^* C^\alpha - C C^{*\alpha} ) + ( \Phi^{*\alpha\beta} \Phi_\beta - \Phi^{\alpha\beta} \Phi_\beta^* ) . \quad (7.8b)$$

For a while let us concentrate on certain useful and auxiliary formulas and relationships. In the first place, one can notice that eq. (7.8) may be considered as another form for reading eq. (7.5). In other words, the following equality

$$\nabla_\alpha (\bar{\Psi} \Gamma^\alpha \Psi) = \bar{\Psi} (\overleftarrow{\partial}_\alpha - B_\alpha) \Gamma^\alpha \Psi + \bar{\Psi} \Gamma^\alpha (\overrightarrow{\partial}_\alpha + B_\alpha) \Psi \quad (7.9a)$$

holds. With the notation

$$(\overrightarrow{\nabla}_\alpha + B_\alpha) \Psi = \overrightarrow{D}_\alpha \Psi, \quad \bar{\Psi} (\overleftarrow{\nabla}_\alpha - B_\alpha) = \bar{\Psi} \overleftarrow{D}_\alpha \quad (7.9b)$$

eq. (7.9a) reads as

$$\nabla_\alpha (\bar{\Psi} \Gamma^\alpha \Psi) = \bar{\Psi} \overleftarrow{D}_\alpha \Gamma^\alpha \Psi + \bar{\Psi} \Gamma^\alpha \overrightarrow{D}_\alpha \Psi . \quad (7.9c)$$

It is particularly remarkable that the formula (7.9c) provides us with a hint about a general rule for covariant  $\nabla_\alpha$  differentiating any bilinear combinations of  $\Psi$  and  $\bar{\Psi}$ . As prescribed by (7.9c), instead of the action of  $\nabla_\alpha$  one may act by  $\overleftarrow{D}_\alpha$  and by  $\overrightarrow{D}_\alpha$  from the right and from the left respectively, but not observing a generally covariant vector characteristic of the matrix  $\Gamma^\alpha(x)$  in the middle.

What is more, such a rule will work always at any complicated bilinear functions. For instance, let us consider an expression

$$\nabla^\alpha [ \bar{\Psi} \Gamma^\rho(x) \Gamma^\sigma(x) \Xi ] .$$

In accordance with defining properties of the covariant derivative we can proceed

$$\begin{aligned} \nabla^\alpha [ \bar{\Psi} \Gamma^\rho \Gamma^\sigma \Xi ] &= \\ &= (\partial_\alpha \bar{\Psi}) \Gamma^\rho \Gamma^\sigma \Xi + \bar{\Psi} \Gamma^\rho_{;\alpha} \Gamma^\sigma \Xi + \bar{\Psi} \Gamma^\rho \Gamma^\sigma_{;\alpha} \Xi + \bar{\Psi} \Gamma^\rho \Gamma^\sigma (\partial_\alpha \Xi) . \end{aligned}$$

Hanging the covariant derivatives of  $\Gamma$ -matrices with commutator as shown in (7.7), we obtain

$$\begin{aligned} \nabla^\alpha [ \bar{\Psi} \Gamma^\rho \Gamma^\sigma \Xi ] &= \\ &= (\partial_\alpha \bar{\Psi}) \Gamma^\rho \Gamma^\sigma \Xi + \bar{\Psi} (\Gamma^\rho B_\alpha - B_\alpha \Gamma^\rho) \Gamma^\sigma \Xi + \bar{\Psi} \Gamma^\rho (\Gamma^\sigma B_\alpha - B_\alpha \Gamma^\sigma) \Xi + \bar{\Psi} \Gamma^\rho \Gamma^\sigma (\partial_\alpha \Xi) . \end{aligned}$$

Two terms containing  $\Gamma^\rho B_\alpha \Gamma^\sigma$  cancel each other, and the remaining can be rewritten as

$$\nabla^\alpha ( \bar{\Psi} \Gamma^\rho \Gamma^\sigma \Xi ) = \bar{\Psi} \overleftarrow{D}_\alpha \Gamma^\rho \Gamma^\sigma \Xi + \bar{\Psi} \Gamma^\rho \Gamma^\sigma \overrightarrow{D}_\alpha \Xi . \quad (7.10a)$$

Eq. (7.10a) is equivalent to

$$\nabla_\alpha (\Gamma^\rho \Gamma^\sigma) = -B_\alpha \Gamma^\rho \Gamma^\sigma + \Gamma^\rho \Gamma^\sigma B_\alpha . \quad (7.10b)$$

An analogous formula for a bilinear combination with any tensor structure follows immediately by induction. Really, let  $\Gamma^{(n)}$  be  $\Gamma^{(n)} = \Gamma^{\rho_1} \Gamma^{\rho_2} \dots \Gamma^{\rho_n}$ , then

$$\nabla_\alpha \Gamma^{(n)} = \Gamma^{(n)} B_\alpha - B_\alpha \Gamma^{(n)} ; \quad (7.11a)$$

so that for  $\Gamma^{(n+1)} = \Gamma^{(n)} \Gamma^\rho$  we can easily derive

$$\begin{aligned} \nabla_\alpha \Gamma^{(n+1)} &= \nabla_\alpha [\Gamma^{(n)} \Gamma^\rho] = [\nabla_\alpha \Gamma^{(n)}] \Gamma^\rho + \Gamma^{(n)} [\nabla_\alpha \Gamma^\rho] = \\ &= [\Gamma^{(n)} B_\alpha - B_\alpha \Gamma^{(n)}] \Gamma^\rho + \Gamma^{(n)} [\Gamma^\rho B_\alpha - B_\alpha \Gamma^\rho] = \Gamma^{(n)} \Gamma^\rho B_\alpha - B_\alpha \Gamma^{(n)} \Gamma^\rho , \end{aligned}$$

which is what we need

$$\nabla_\alpha \Gamma^{(n+1)} = -B_\alpha \Gamma^{(n+1)} + \Gamma^{(n+1)} B_\alpha . \quad (7.11b)$$

From formulas (7.11) one can produce the commutation relations

$$\Gamma^{(n)} \overrightarrow{D}_\sigma = \overrightarrow{D}_\sigma \Gamma^{(n)} , \quad \Gamma^{(n)} \overleftarrow{D}_\sigma = \overleftarrow{D}_\sigma \Gamma^{(n)} . \quad (7.12)$$

Now we are ready to write down a Lagrange function that yields eqs. (7.2) and (7.4). At this, two formulas will be taken into account (see 7.12)):

$$\Gamma^\rho(\nabla_\sigma + B_\sigma) = (\nabla_\sigma + B_\sigma)\Gamma^\rho, \quad \Gamma^\rho(\overleftarrow{\nabla}_\sigma - B_\sigma) = (\overleftarrow{\nabla}_\sigma - B_\sigma)\Gamma^\rho, \quad (7.13a)$$

or in a short form

$$\Gamma^\rho \overrightarrow{D}_\sigma = \overrightarrow{D}_\sigma \Gamma^\rho, \quad \Gamma^\rho \overleftarrow{D}_\sigma = \overleftarrow{D}_\sigma \Gamma^\rho. \quad (7.13b)$$

Bearing in mind that the basic equations for  $\Psi$  and  $\bar{\Psi}$  look as

$$[\Gamma^\alpha \overrightarrow{D}_\alpha - ]\Psi = 0, \quad (7.14a)$$

$$\bar{\Psi}(\Gamma^\alpha \overleftarrow{D}_\alpha + m) = 0, \quad (7.14b)$$

a generally covariant Lagrangian for this system can be chosen in the form

$$L = +\frac{1}{2} [\bar{\Psi} \Gamma^\rho \overrightarrow{D}_\rho \Psi - \bar{\Psi} \Gamma^\rho \overleftarrow{D}_\rho \Psi] - m \bar{\Psi} \Psi. \quad (7.15a)$$

In terms of tensor components, the  $L$  will take on the form

$$\begin{aligned} L = & \frac{1}{2} [-\sigma (C^{*\alpha} \nabla_\alpha C + C^\alpha \nabla_\alpha C^*) + \sigma (C^* \nabla_\alpha C^\alpha + C \nabla_\alpha C^{*\alpha}) + \\ & + (\Phi^{*\alpha\beta} \nabla_\alpha \Phi_\beta + \Phi^{\alpha\beta} \nabla_\alpha \Phi_\beta^*) + (\Phi_\alpha^* \nabla_\beta \Phi^{\alpha\beta} + \Phi_\alpha \nabla_\beta \Phi^{*\alpha\beta})] - \\ & - m [\sigma C^* C + C_\alpha^* C^\alpha - (\Phi_\alpha^* C^\alpha + \Phi_\alpha C^{*\alpha}) + \frac{1}{2} \Phi_{\alpha\beta}^* \Phi^{\alpha\beta}]. \end{aligned} \quad (7.15b)$$

The limiting case of a massless particle follows from (7.15) by a formal change (similar to the flat space-time)

$$m(\bar{\Psi} \Psi) \implies \bar{\Psi} P \Psi, \quad P = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & I & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & I \end{pmatrix}, \quad (7.16a)$$

which transforms the last term in (7.15b)

$$\begin{aligned} -m [\sigma C^* C + C_\alpha^* C^\alpha - (\Phi_\alpha^* C^\alpha + \Phi_\alpha C^{*\alpha}) + \frac{1}{2} \Phi_{\alpha\beta}^* \Phi^{\alpha\beta}] \implies \\ - [\sigma C^* C + C_\alpha^* C^\alpha + \frac{1}{2} \Phi_{\alpha\beta}^* \Phi^{\alpha\beta}]. \end{aligned} \quad (7.16b)$$

## 8. Massless equation and conformal invariance

Section 8 deals with a property of conformal invariance as it looks for 15-component theory of massless spin 1 particle. An original equation is

$$[\Gamma^\alpha (\partial_\alpha + B_\alpha) - P] \Psi = 0, \quad (8.1)$$

projective matrix  $P$  was defined by (7.16a). Let the metric tensors of two curved space-time models differ by a factor - function:

$$\tilde{g}_{\alpha\beta}(x) = \varphi^2(x) g_{\alpha\beta}(x). \quad (8.2)$$

Correspondingly, we are to use two vierbeins connected with each other by means of the same factor  $\varphi$ :

$$\tilde{e}_{(a)}^\alpha = \varphi^{-1} e_{(a)}^\alpha, \quad \tilde{e}_{(a)\alpha} = \varphi e_{(a)\alpha}. \quad (8.3)$$

For two sets of Christoffel symbols we have

$$\begin{aligned} \tilde{\Gamma}_{\alpha\beta,\rho} &= \frac{1}{2} [\partial_\alpha \tilde{g}_{\beta\rho} + \partial_\beta \tilde{g}_{\alpha\rho} - \partial_\rho \tilde{g}_{\alpha\beta}] = \\ &= \varphi^2 \Gamma_{\alpha\beta,\rho} + \varphi [(\partial_\alpha \varphi) g_{\beta\rho} + (\partial_\beta \varphi) g_{\alpha\rho} - (\partial_\rho \varphi) g_{\alpha\beta}], \end{aligned}$$

or, with the use of the representative  $\tilde{\Gamma}_{\alpha\beta}^\sigma = \tilde{g}^{\sigma\rho}\Gamma_{\alpha\beta,\rho}$ , in another form

$$\tilde{\Gamma}_{\alpha\beta}^\sigma = \Gamma_{\alpha\beta}^\sigma + \frac{1}{\varphi} [(\partial_\alpha \varphi) \delta_\beta^\sigma + (\partial_\beta \varphi) \delta_\alpha^\sigma - g^{\sigma\rho} (\partial_\rho \varphi) g_{\alpha\beta}] . \quad (8.4)$$

Let us compare two sets of the Ricci rotation coefficients:

$$\begin{aligned} \tilde{\gamma}_{abc}(x) &= -[ \partial_\beta \tilde{e}_{(a)\alpha} - \tilde{\Gamma}_{\beta\alpha}^\rho \tilde{e}_{(a)\rho} ] \tilde{e}_{(b)}^\alpha \tilde{e}_{(c)}^\beta = \\ &= -[ \partial_\alpha (\varphi e_{(a)\alpha}) - \Gamma_{\beta\alpha}^\rho \varphi e_{(a)\rho} - \\ &\quad - \frac{1}{\varphi} \left( (\partial_\beta \varphi) \delta_\alpha^\rho + (\partial_\alpha \varphi) \delta_\beta^\rho - g^{\rho\sigma} (\partial_\sigma \varphi) g_{\beta\alpha} \right) \varphi e_{(a)\rho} ] \frac{1}{\varphi^2} e_{(b)}^\alpha e_{(c)}^\beta . \end{aligned}$$

and further

$$\tilde{\gamma}_{abc} = -[ \varphi e_{(a)\alpha;\beta} - \frac{\partial \varphi}{\partial x^\alpha} e_{(a)\beta} + e_{(a)}^\sigma \frac{\partial \varphi}{\partial x^\sigma} g_{\alpha\beta} ] \frac{1}{\varphi^2} e_{(b)}^\alpha e_{(c)}^\beta .$$

So, the formula we need is

$$\tilde{\gamma}_{abc} = \frac{1}{\varphi} \gamma_{abc} + \frac{1}{\varphi^2} (\partial_\sigma \varphi) [e_{(b)}^\sigma g_{ac} - e_{(a)}^\sigma g_{bc}] . \quad (8.5)$$

An equation (8.1) in a space-time with the metric tensor  $\tilde{g}_{\alpha\beta}(x)$  is

$$[ \tilde{\Gamma}^\alpha (\partial_\alpha + \tilde{B}_\alpha) - P ] \tilde{\Psi} = 0 , \quad (8.6)$$

or

$$[ \Gamma^c (\tilde{e}_{(c)}^\alpha \partial_\alpha + \frac{1}{2} J^{ab} \tilde{\gamma}_{abc}) - P ] \tilde{\Psi} = 0 . \quad (8.7)$$

Substituting expressions for a new tetrad and new Ricci coefficients (tilded) in terms of old (not-tilded) ones, we get to

$$[\Gamma^c (e_{(c)}^\sigma \partial_\sigma + \frac{1}{2} J^{ab} \gamma_{abc}) + \frac{1}{2} \Gamma^c J^{ab} [ (e_{(b)}^\sigma g_{ac} - e_{(a)}^\sigma g_{bc}) (\varphi^{-1} \partial_\sigma \varphi) ] \tilde{\Psi} = \varphi P \tilde{\Psi} .$$

and further

$$[ \Gamma^\sigma (\partial_\sigma + B_\sigma) + \Gamma_a J^{ab} e_{(b)}^\sigma (\varphi^{-1} \partial_\sigma \varphi) ] \tilde{\Psi} = \varphi P \tilde{\Psi} . \quad (8.8)$$

With the use of block-representations for all matrices, consider a combination  $\Gamma_a J^{ab}$ :

$$\Gamma_a J^{ab} = \begin{pmatrix} 0 & G_a V^{ab} & 0 & 0 \\ 0 & 0 & 0 & K_a (V \otimes V)^{ab} \\ 0 & 0 & 0 & K_a (V \otimes V)^{ab} \\ 0 & 0 & \Lambda_a V^{ab} & 0 \end{pmatrix} . \quad (8.9)$$

Allowing for relations

$$\begin{aligned} (G_a)_{(0)}^l (V^{ab})_l{}^k &= +3g^{bk} , \quad (\Lambda_a)_{ps}^l (V^{ab})_l{}^k = \delta_{ps}^{bk} , \\ (K_a)_l{}^{mn} [(V \otimes V)^{ab}]_{mn}{}^{ps} &= 2(-g^{bp} \delta_l^s + g^{bs} \delta_l^p) , \end{aligned}$$

an expression for  $\Gamma_a J^{ab}$  is led to

$$\Gamma_a J^{ab} = \begin{pmatrix} 0 & +3g^{bk} & 0 & 0 \\ 0 & 0 & 0 & 2(-g^{bp} \delta_l^s + g^{bs} \delta_l^p) \\ 0 & 0 & 0 & 2(-g^{bp} \delta_l^s + g^{bs} \delta_l^p) \\ 0 & 0 & \delta_{ps}^{bk} & 0 \end{pmatrix} . \quad (8.10)$$

From this, with taking in mind eq. (2.12), we find

$$\Gamma_a J^{ab} = \begin{pmatrix} 0 & 3(G^b)_{(0)}^k & 0 & 0 \\ 0 & 0 & 0 & 2(K^b)_l{}^{ps} \\ 0 & 0 & 0 & 2(K^b)_l{}^{ps} \\ 0 & 0 & (\Lambda^b)_{ps}^k & 0 \end{pmatrix} = \Gamma^b \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & +3I & 0 & 0 \\ 0 & 0 & +I & 0 \\ 0 & 0 & 0 & +2I \end{pmatrix} . \quad (8.11a)$$



In the end, we arrive at

$$\Gamma_a J^{ab} = \Gamma^b (+3P'_4 + P_4 + 2P_6) , \quad (8.11b)$$

where a special notation for projective operators on 4-dimensional and 6-dimensional subspaces is used. Hence, eq. (8.9) reads as

$$\Gamma^\sigma [ \partial_\sigma + B_\sigma + (3P'_4 + P_4 + 2P_6)(\varphi^{-1} \partial_\sigma \varphi) ] \tilde{\Psi} = \varphi P \tilde{\Psi} . \quad (8.12)$$

With the use of block-forms of all matrices involved in (8.12) and taking the notation

$$\tilde{\Psi} = (\tilde{\Phi}, \tilde{C}, \tilde{A}, \tilde{F}) , \quad (8.13)$$

eq. (8.12) is shown to be equivalent to a set (compare with (6.21))

$$G^\alpha [\partial_\alpha + V_\alpha + 3\partial_\alpha \ln \varphi] \tilde{C} = \varphi \tilde{\Phi} , \quad (8.14a)$$

$$K^\alpha [\partial_\alpha + (V \otimes V)_\alpha + 2\partial_\alpha \ln \varphi] \tilde{F} = \varphi \tilde{C} , \quad (8.14b)$$

$$\sigma \Delta^\alpha \partial_\alpha \tilde{\Phi} + K^\alpha [\partial_\alpha + (V \otimes V)_\alpha + 2\partial_\alpha \ln \varphi] \tilde{F} = 0 , \quad (8.14c)$$

$$\Lambda^\alpha [\partial_\alpha + V_\alpha + \partial_\alpha \ln \varphi] \tilde{A} = \varphi \tilde{F} \quad (8.14d)$$

Let us find out how will look these equations after substitution the field components in the form

$$\tilde{A}_l = \varphi^{-1} A_l , \tilde{F}_{mn} = \varphi^{-2} F_{mn} , \tilde{C}_l = \varphi^{-3} C_l , \tilde{\Phi} = \varphi^{-4} \Phi . \quad (8.15)$$

It is straightforward to obtain

$$G^\alpha [\partial_\alpha + V_\alpha] C = \Phi , \quad (8.16a)$$

$$K^\alpha [\partial_\alpha + (V \otimes V)_\alpha] F = C , \quad (8.16b)$$

$$\varphi^2 \sigma \Delta^\alpha \partial_\alpha \tilde{\Phi} + K^\alpha [\partial_\alpha + (V \otimes V)_\alpha] F = 0 , \quad (8.16c)$$

$$\Lambda^\alpha [\partial_\alpha + V_\alpha] A = F . \quad (8.16d)$$

Take notice that apparently eq. (8.16c) does not coincides with a respective equation in  $g_{\alpha\beta}$ -space. However, if the vanishing of supplementary scalar and vector components  $\tilde{\Phi} = 0$  ,  $\tilde{C} = 0$  is taken into consideration, then effectively remaining two equations in (8.16) will read as

$$K^\alpha [\partial_\alpha + (V \otimes V)_\alpha] F = 0 ,$$

$$\Lambda^\alpha [\partial_\alpha + V_\alpha(x)] A = F , \quad (8.17)$$

which are precisely required ones for a set  $(\Phi = 0, C = 0, A, F)$  in  $g_{\alpha\beta}$ -space. This means that the property of conformal invariance holds.

To avoid misleading it should be mentioned that just accomplished study of conformal invariance in 15-component theory of a massless vector particle does not add much to yet known for a conventional 10-component theory; rather it lies in the freedom from any internal conflict in the theory. This is what should be expected because of equivalence demonstrated above between 15- and 10-component models in a free particle case.

## 9. Canonical energy-momentum tensor

In Sec. 9 we are going to study a question about energy-momentum tensor for a generalized spin 1 particle. At this it is convenient to exploit a matrix formalism, when two basic equations are given by (for much generality, an external electromagnetic field will be taken into account)

$$(\Gamma^\alpha \vec{D}_\alpha - m) \Psi = 0 , \quad \bar{\Psi} (\Gamma^\alpha \overleftarrow{D}_\alpha + m) = 0 , \quad (9.1a)$$

where

$$\vec{D}_\alpha = \vec{\nabla}_\alpha + B_\alpha - ig A_\alpha , \quad \overleftarrow{D}_\alpha = \overleftarrow{\nabla}_\alpha - B_\alpha + ig A_\alpha , \quad g \equiv e/\hbar c . \quad (9.1b)$$

Let a tensor quantity  $W_\beta{}^\alpha$  be

$$W_\beta{}^\alpha = \bar{\Psi} \Gamma^\alpha \vec{D}_\beta \Psi = \bar{\Psi} \Gamma^\alpha (\partial_\beta + B_\beta) \Psi - ig A_\beta (\bar{\Psi} \Gamma^\alpha \Psi) . \quad (9.2)$$

The aim is to calculate a divergence over it. To this end, act on the first equation in (9.1) from the left by an operator  $\bar{\Psi} \vec{D}_\beta$ , and multiply the second from the right by  $\text{na } \vec{D}_\beta \Psi$ , and sum results, then we have

$$\bar{\Psi} \vec{D}_\beta \Gamma^\alpha \vec{D}_\alpha \Psi + \bar{\Psi} \overleftarrow{D}_\alpha \Gamma^\alpha \vec{D}_\beta \Psi = 0, \quad (9.3a)$$

an further

$$\bar{\Psi} \Gamma^\alpha [(\vec{D}_\beta \vec{D}_\alpha - \vec{D}_\alpha \vec{D}_\beta) + \overleftarrow{D}_\alpha \vec{D}_\beta] \Psi + \bar{\Psi} \overleftarrow{D}_\alpha \Gamma^\alpha \vec{D}_\beta \Psi = 0. \quad (9.3b)$$

Now, placing a term with commutator  $[\vec{D}_\beta, \vec{D}_\alpha]_-$  on the right, we get to

$$\bar{\Psi}(\Gamma^\alpha \vec{D}_\alpha + \overleftarrow{D}_\alpha \Gamma^\alpha) \vec{D}_\beta \Psi = \bar{\Psi} \Gamma^\alpha [\vec{D}_\alpha, \vec{D}_\beta]_- \Psi. \quad (9.3c)$$

Let us show that an expression on the left can be thought as a pure divergence of  $W_\beta^\alpha$ . We are to start with

$$\begin{aligned} & \bar{\Psi}(\Gamma^\alpha \vec{D}_\alpha + \overleftarrow{D}_\alpha \Gamma^\alpha) \vec{D}_\beta \Psi = \\ & = \bar{\Psi}[\Gamma^\alpha(\vec{\nabla}_\alpha + B_\alpha - igA_\alpha) + (\overleftarrow{\nabla}_\alpha - B_\alpha + igA_\alpha)\Gamma^\alpha] \vec{D}_\beta \Psi = \\ & = \bar{\Psi}[\Gamma^\alpha \vec{\nabla}_\alpha + (\Gamma^\alpha B_\alpha - B_\alpha \Gamma^\alpha) + \overleftarrow{\nabla}_\alpha \Gamma^\alpha] \vec{D}_\beta \Psi. \end{aligned} \quad (9.4a)$$

Taking into consideration eq. (7.7), relation (9.4a) can be transformed into

$$\begin{aligned} & \bar{\Psi}(\Gamma^\alpha \vec{D}_\alpha + \overleftarrow{D}_\alpha \Gamma^\alpha) \vec{D}_\beta \Psi = \\ & = \bar{\Psi}(\overleftarrow{\nabla}_\alpha \Gamma^\alpha + \Gamma^\alpha_{;\alpha} + \Gamma^\alpha \overleftarrow{\nabla}_\alpha) \vec{D}_\beta \Psi = \nabla_\alpha (\bar{\Psi} \Gamma^\alpha \vec{D}_\beta \Psi). \end{aligned} \quad (9.4b)$$

Hence, eq. (9.3c) is equivalent to

$$\nabla_\alpha [W_\beta^\alpha] = \bar{\Psi} \Gamma^\alpha [\vec{D}_\alpha, \vec{D}_\beta]_- \Psi. \quad (9.5)$$

Let us consider in more detail the commutator

$$[\vec{D}_\alpha, \vec{D}_\beta]_- = -igF_{\alpha\beta} + D_{\alpha\beta}, \quad (9.6a)$$

where

$$F_{\alpha\beta} = \frac{\partial A_\beta}{\partial x^\alpha} - \frac{\partial A_\alpha}{\partial x^\beta}, \quad D_{\alpha\beta} = \frac{\partial B_\beta}{\partial x^\alpha} - \frac{\partial B_\alpha}{\partial x^\beta} + (B_\alpha B_\beta - B_\beta B_\alpha). \quad (9.6b)$$

For first term on the right in expression for  $D_{\alpha\beta}$ , it is easily to obtain a representation

$$\begin{aligned} & \partial_\alpha B_\beta - \partial_\beta B_\alpha = \nabla_\alpha B_\beta - \nabla_\beta B_\alpha = \\ & = \frac{1}{2} J^{ab} \nabla_\alpha (e_{(a)}^\nu e_{(b)\nu;\beta}) - \frac{1}{2} J^{ab} \nabla_\beta (e_{(a)}^\nu e_{(b)\nu;\alpha}) = \\ & = \frac{1}{2} J^{ab} e_{(a)}^\nu [e_{(b)\nu;\beta;\alpha} - e_{(b)\nu;\alpha;\beta}] + \frac{1}{2} J^{ab} [e_{(a)\nu;\alpha} e_{(b);\beta}^\nu - e_{(a)\nu;\beta} e_{(b);\alpha}^\nu]. \end{aligned} \quad (9.7)$$

For second one it follows

$$\begin{aligned} & (B_\alpha B_\beta - B_\beta B_\alpha) = \\ & = (\frac{1}{2} J^{ab} e_{(a)}^\nu e_{(b)\nu;\alpha}) (\frac{1}{2} J^{kl} e_{(k)}^\mu e_{(l)\mu;\beta}) - (\frac{1}{2} J^{kl} e_{(k)}^\mu e_{(l)\mu;\beta}) (\frac{1}{2} J^{ab} e_{(a)}^\nu e_{(b)\nu;\alpha}) = \\ & = \frac{1}{4} (J^{ab} J^{kl} - J^{kl} J^{ab}) \left[ (e_{(a)}^\nu e_{(b)\nu;\alpha}) e_{(k)}^\mu e_{(l)\mu;\beta} \right]. \end{aligned}$$

Now, with the use of commutation relations (see (6.11))

$$[J^{ab}, J^{kl}]_- = (-J^{kb} g^{la} + J^{lb} g^{ka}) - (-J^{ka} g^{lb} + J^{la} g^{kb}),$$

we get

$$(B_\alpha B_\beta - B_\beta B_\alpha) = -\frac{1}{2} J^{ab} [e_{(a)\nu;\alpha} e_{(b);\beta}^\nu - e_{(a)\nu;\beta} e_{(b);\alpha}^\nu]. \quad (9.8)$$

Accounting for eqs. (9.7) and (9.8), expression for  $D_{\alpha\beta}$  can be led to the form

$$\begin{aligned} D_{\alpha\beta} &= \frac{1}{2} J^{ab} e_{(a)}^\nu [ e_{(b)\nu;\beta;\alpha} - e_{(b)\nu;\alpha;\beta} ] = \\ &= \frac{1}{2} J^{ab} e_{(a)}^\nu [ e_{(b)}^\rho R_{\rho\nu\beta\alpha}(x) ] = \frac{1}{2} J^{\nu\rho}(x) R_{\nu\rho\alpha\beta}(x) , \end{aligned} \quad (9.9)$$

where  $R_{\nu\rho\alpha\beta}(x)$  designates a Riemann curvature tensor. Thus, eq. (9.5) reads as

$$\nabla_\alpha W_\beta{}^\alpha = -ig J^\alpha F_{\alpha\beta} + \frac{1}{2} R_{\nu\rho\alpha\beta} \bar{\Psi} \Gamma^\alpha J^{\nu\rho} \Psi . \quad (9.10)$$

It is useful to perform some transformation over second term on the right:

$$\begin{aligned} &\frac{1}{2} R_{\nu\rho\alpha\beta} \bar{\Psi} \Gamma^\alpha J^{\nu\rho} \Psi = \\ &= \frac{1}{2} R_{\nu\rho\alpha\beta} \bar{\Psi} \left( \frac{1}{2} [ \Gamma^\alpha J^{\nu\rho} - J^{\nu\rho} \Gamma^\alpha ] + \frac{1}{2} [ \Gamma^\alpha J^{\nu\rho} + J^{\nu\rho} \Gamma^\alpha ] \right) \Psi . \end{aligned}$$

Further, with the use of commutation relation (see (7.6))

$$\Gamma^\alpha J^{\nu\rho} - J^{\nu\rho} \Gamma^\alpha = g^{\alpha\nu}(x) \Gamma^\rho - g^{\alpha\rho}(x) \Gamma^\nu ,$$

one can produce

$$\begin{aligned} &\frac{1}{2} R_{\nu\rho\alpha\beta} \bar{\Psi} \Gamma^\alpha J^{\nu\rho} \Psi = \\ &= \frac{1}{2} R_{\alpha\beta} J^\alpha + \frac{1}{4} R_{\nu\rho\alpha\beta} \bar{\Psi} (\Gamma^\alpha J^{\nu\rho} + J^{\nu\rho} \Gamma^\alpha) \Psi . \end{aligned} \quad (9.11)$$

Correspondingly, eq. (9.10) takes on the form

$$\nabla_\alpha [ W_\beta{}^\alpha ] = -ig J^\alpha F_{\alpha\beta} + \frac{1}{2} J^\alpha R_{\alpha\beta} + \frac{1}{4} R_{\nu\rho\alpha\beta} \bar{\Psi} (\Gamma^\alpha J^{\nu\rho} + J^{\nu\rho} \Gamma^\alpha) \Psi . \quad (9.12a)$$

This relation can be rewritten as

$$\nabla_\alpha [ W_\beta{}^\alpha ] = J^\alpha ( -ig F_{\alpha\beta} + \frac{1}{2} R_{\alpha\beta} ) + \frac{1}{4} R_{\nu\rho\alpha\beta} \bar{\Psi} (\Gamma^\alpha J^{\nu\rho} + J^{\nu\rho} \Gamma^\alpha) \Psi , \quad (9.12b)$$

which generalizes a known formula established by V.A. Fock [49, eq. (56)] at studying a spin 1/2 particle on the background of a curved space-time model. A single formal difference consists in occurrence of one additional term proportional to Riemann curvature tensor on the left.

It may be checked quite easily that, in the established formula (9.12b) being applied to spin 1/2 particle case, this additional  $R$ -dependent term will vanish identically. To this end, let us consider more closely a combination of Dirac matrices

$$(\gamma^\alpha j^{\nu\rho} + j^{\nu\rho} \gamma^\alpha) = \frac{1}{4} [ \gamma^\alpha (\gamma^\nu \gamma^\rho - \gamma^\rho \gamma^\nu) + (\gamma^\nu \gamma^\rho - \gamma^\rho \gamma^\nu) \gamma^\alpha ]$$

Multiplying the known formula for the product of the ordinary Dirac matrices [12]

$$\gamma^a \gamma^b \gamma^c = \gamma^a g^{bc} - \gamma^b g^{ac} + \gamma^c g^{ab} + i\gamma^5 \epsilon^{abcd} \gamma_d \quad (9.13a)$$

by the following tetrad construction  $e_{(a)}^\alpha e_{(b)}^\beta e_{(c)}^\rho$ , one obtains

$$\begin{aligned} \gamma^\alpha(x) \gamma^\beta(x) \gamma^\rho(x) &= [ \gamma^\alpha(x) g^{\beta\rho}(x) - \gamma^\beta(x) g^{\alpha\rho}(x) + \\ &+ \gamma^\rho(x) g^{\alpha\beta}(x) + i\gamma^5 \epsilon^{\alpha\beta\rho\sigma}(x) \gamma_\sigma(x) ] . \end{aligned} \quad (9.13b)$$

Here a generally covariant Levi-Civita symbol is given by

$$\epsilon^{\alpha\beta\rho\sigma}(x) = e_{(a)}^\alpha e_{(b)}^\beta e_{(c)}^\rho e_{(d)}^\sigma \epsilon^{abcd} . \quad (9.13c)$$

Now, with the use of (9.13b), one can produce

$$\gamma^\alpha(x)j^{\nu\rho}(x) + j^{\nu\rho}(x)\gamma^\alpha(x) = i\gamma^5 \epsilon^{\alpha\nu\rho\sigma}(x) \gamma_\sigma(x). \quad (9.14a)$$

So, we have arrived to

$$\frac{1}{4} R_{\nu\rho\alpha\beta}(x) \bar{\Psi}(\Gamma^\alpha J^{\nu\rho} + J^{\nu\rho}\Gamma^\alpha)\Psi = \frac{1}{4} R_{\nu\rho\alpha\beta}(x) \epsilon^{\alpha\nu\rho\sigma}(x) \bar{\Psi}\gamma^5 \gamma_\sigma(x)\Psi \equiv 0; \quad (9.14)$$

where it is taken into account the known symmetry of the curvature tensor under cyclic permutation over any three indices so that convolution of  $R_{\dots}$  with  $\epsilon_{\dots}$  over three indices equals to zero.

Returning again to (9.12b), with the use of the properties of invariant-form matrix  $\eta$  (see (3.3) and (7.1b)):

$$\eta^{-1}[\Gamma^\alpha(x)]^+\eta = -\Gamma^\alpha(x), \quad \eta^{-1}[J^{\nu\rho}(x)]^+\eta = -J^{\nu\rho}(x), \quad (9.15)$$

one can find that on the right in (9.12a) the first and the third terms are real-valued, whereas the second is imaginary:

$$\begin{aligned} \nabla_\alpha W_\beta{}^\alpha &= Re(x) + i Im(x), \quad Im(x) = -\frac{i}{2} J^\alpha R_{\alpha\beta}, \\ Re(x) &= -ig J^\alpha F_{\alpha\beta} + \frac{1}{4} R_{\nu\rho\alpha\beta} \bar{\Psi}(\Gamma^\alpha J^{\nu\rho} + J^{\nu\rho}\Gamma^\alpha)\Psi. \end{aligned} \quad (9.16)$$

Now, we need to isolate real and imaginary parts on the left in (9.12a) too. To this end, let us find a complex conjugate tensor  $(W_\beta^\alpha)^*$ :

$$\begin{aligned} (W_\beta^\alpha)^+ &= [\Psi^+ \eta \Gamma^\alpha (\nabla_\beta + B_\beta - ig A_\beta) \Psi]^+ = \\ &= -\bar{\Psi} \Gamma^\alpha (\overleftarrow{\nabla}_\beta - B_\beta + ig A_\beta) \Psi = -\bar{\Psi} \overleftarrow{D}_\beta \Gamma^\alpha \Psi. \end{aligned} \quad (9.17)$$

With the notation

$$\begin{aligned} Re(W_\beta^\alpha) &= \frac{1}{2} [W_\beta^\alpha + (W_\beta^\alpha)^+] = T_\beta^\alpha, \\ Im(W_\beta^\alpha) &= \frac{1}{2i} [W_\beta^\alpha - (W_\beta^\alpha)^+] = U_\beta^\alpha, \end{aligned} \quad (9.18)$$

eq. (9.16) will split into two real-valued ones

$$\nabla_\alpha (T_\beta^\alpha) = -ig J^\alpha(x) F_{\alpha\beta} + \frac{1}{4} R_{\nu\rho\alpha\beta} \bar{\Psi} (\Gamma^\alpha J^{\nu\rho} + J^{\nu\rho} \Gamma^\alpha) \Psi. \quad (9.19)$$

$$\nabla_\alpha (U_\beta^\alpha) = -\frac{i}{2} J^\alpha R_{\alpha\beta}, \quad (9.20)$$

As it is readily checked, eq. (9.20) represent in essence a direct consequence of the conserved current law. Really, in accordance with definition for  $U_\beta^\alpha$  we have

$$\begin{aligned} U_\beta^\alpha &= \frac{1}{2i} [\bar{\Psi} \Gamma^\alpha (\partial_\beta + B_\beta - iA_\beta) \Psi + \bar{\Psi} (\overleftarrow{\partial}_\beta - B_\beta + iA_\beta) \Gamma^\alpha \Psi] = \\ &= \frac{1}{2i} \nabla_\beta (\bar{\Psi} \Gamma^\alpha \Psi) = \frac{1}{2i} \nabla_\beta J^\alpha. \end{aligned} \quad (9.21)$$

Therefore, eq. (9.19) can be led to the form

$$\nabla_\alpha \nabla_\beta J^\alpha = R_{\alpha\beta} J^\alpha, \quad (9.22a)$$

and further

$$(\nabla_\alpha \nabla_\beta - \nabla_\beta \nabla_\alpha) J^\alpha + \nabla_\beta \nabla_\alpha J^\alpha = J^\alpha R_{\alpha\beta}.$$

From this, with bearing in mind the conservation law for  $J^\alpha$ , it follows an identity

$$J^\rho(x) R_{\rho\beta\alpha}(x) \equiv J^\alpha(x) R_{\alpha\beta}(x). \quad (9.22b)$$

Thus, eq. (9.20) does not contain anything new in addition to current conservation law. As for eq. (9.19), for  $T_\beta^\alpha$  we have

$$\begin{aligned} T_\beta^\alpha &= \frac{1}{2} [\bar{\Psi} \Gamma^\alpha \vec{D}_\beta \Psi - \bar{\Psi} \Gamma^\alpha \overleftarrow{D}_\beta \Psi] = \\ &= \frac{1}{2} [\bar{\Psi} \Gamma^\alpha (\vec{\nabla}_\beta + B_\beta) \Psi - \bar{\Psi} \Gamma^\alpha (\overleftarrow{\nabla}_\beta - B_\beta) \Psi] - ig J^\alpha A_\beta . \end{aligned} \quad (9.23a)$$

and a conservation law reads as follows

$$\nabla_\alpha (T_\beta^\alpha) = -ig J^\alpha F_{\alpha\beta} + \frac{1}{4} R_{\nu\rho\sigma\beta} \bar{\Psi} (\Gamma^\sigma J^{\nu\rho} + J^{\nu\rho} \Gamma^\sigma) \Psi . \quad (9.23b)$$

To proceed with eq. (9.23b), it is a time to remember some facts about ambiguity in determining any energy-momentum tensor. In the Minkowski space-time such freedom in its determining is described as follows: if  $T_b^a(x)$  obeys an equation

$$\partial_a T_b^a = 0 ,$$

then another tensor

$$\bar{T}_b^a(x) = T_b^a(x) + \partial_c [\Omega_b^{[ac]}(x)] , \quad \text{where } \Omega_b^{[ac]}(x) = -\Omega_b^{[ca]}(x) \quad (9.24a)$$

satisfies the same equation as well

$$\partial_a \bar{T}_b^a = 0$$

Obviously, simultaneous existence of the two conservation laws is insured by an elementary formula

$$\partial_a \partial_c \Omega_b^{[ca]}(x) \equiv 0 . \quad (9.24b)$$

As for a curved space-time model, such an equivalence between tensors  $T_\beta^\alpha(x)$  and  $\bar{T}_\beta^\alpha(x)$  holds as well but in a more complicated manner. Indeed, let two tensors be related to each other by

$$\bar{T}_\beta^\alpha(x) = T_\beta^\alpha(x) + \nabla_\rho [\Omega_\beta^{[\alpha\rho]}(x)] . \quad (9.25)$$

Acting on both sides by operation of covariant differentiation  $\nabla_\alpha$ , one produces

$$\nabla_\alpha \bar{T}_\beta^\alpha(x) = \nabla_\alpha T_\beta^\alpha(x) + \nabla_\alpha [\nabla_\rho \Omega_\beta^{[\alpha\rho]}(x)] . \quad (9.26)$$

Now, bearing in mind symmetry properties of the curvature tensor, one obtains

$$\begin{aligned} \nabla_\alpha [\nabla_\rho \Omega_\beta^{[\alpha\rho]}(x)] &= \frac{1}{2} [R_{\alpha\rho\beta}{}^\sigma \Omega_\sigma^{[\alpha\rho]} + R_{\alpha\rho}{}^\alpha{}_\sigma \Omega_\beta^{[\sigma\rho]} + R_{\alpha\rho}{}^\rho{}_\sigma \Omega_\beta^{[\alpha\sigma]}] = \\ &= \frac{1}{2} [R^{\beta\sigma}{}_{\alpha\rho} \Omega_\sigma^{[\alpha\rho]} - R_{\rho\sigma} \Omega_\beta^{[\sigma\rho]} - R_{\alpha\sigma} \Omega_\beta^{[\alpha\sigma]}] , \end{aligned}$$

thus

$$\nabla_\alpha [\nabla_\rho \Omega_\beta^{[\alpha\rho]}(x)] = \frac{1}{2} R_{\beta\sigma\nu\rho} \Omega^{\sigma[\nu\rho]} . \quad (9.27)$$

In the end, eq. (9.26) reads as

$$\nabla_\alpha \bar{T}_\beta^\alpha(x) = \nabla_\alpha T_\beta^\alpha(x) + \frac{1}{2} R_{\beta\sigma\nu\rho} \Omega^{\sigma[\nu\rho]}(x) . \quad (9.28)$$

which, on accounting eq.(9.23b), takes the form

$$\begin{aligned} \nabla_\alpha \bar{T}_\beta^\alpha(x) &= -ig J^\alpha F_{\alpha\beta} + \frac{1}{4} R_{\nu\rho\sigma\beta} \bar{\Psi} (\Gamma^\sigma J^{\nu\rho} + J^{\nu\rho} \Gamma^\sigma) \Psi + \\ &+ \frac{1}{2} R_{\beta\sigma\nu\rho} \Omega^{\sigma[\nu\rho]}(x) . \end{aligned} \quad (9.29a)$$

If the quantity  $\Omega^{\sigma[\nu\rho]}(x)$  is chosen as

$$\Omega^{\sigma[\nu\rho]}(x) = +\frac{1}{2} \bar{\Psi} [\Gamma^\sigma J^{\nu\rho} + J^{\nu\rho} \Gamma^\sigma] \Psi , \quad (9.29b)$$

then second and third terms on the right in (9.29a) cancel each other, and we reach a conservation law

$$\nabla_\alpha \bar{T}_\beta^\alpha(x) = -ig J^\alpha(x) F_{\alpha\beta}(x) . \quad (9.30)$$

## Results

A generalized vector particle theory with the use of an extended set of Lorentz group irreducible representations, including scalar, two 4-vectors, and antisymmetric 2-rang tensor, is investigated both in tensor and in matrix approaches. Initial equations depend upon four complex parameters  $\lambda_i$ , obeying two supplementary conditions, so restriction of the model to the case of electrically neutral vector particle is not a trivial task. A special basis in the space of 15-component wave functions is found where instead of four  $\lambda_i$  only one real-valued quantity  $\sigma$ , a bilinear combination of  $\lambda_i$ , is presented. This  $\lambda$ -parameter is interpreted as an additional electromagnetic characteristic of a charged vector particle, polarizability. It is shown that in this basis  $C$ -operation is reduced to the complex conjugation only, without any accompanying linear transformation. The form of  $C$ -operation in the initial basis is calculated too. Invariant bilinear form matrix in both bases are found and the Lagrange formulation of the whole theory is given. Explicit expressions of the conserved current vector and of the energy-momentum tensor are established. In presence of external electromagnetic fields, two supplementary field components, scalar and 4-vector, give a non-trivial contribution to the Lagrangian and conserved quantities. Restriction to a massless vector particle is determined.

Extension of the whole theory to the case of Riemannian space-time is accomplished. Two methods of obtaining corresponding generally covariant wave equations are elaborated: of tensor- and of tetrad-based ones. Their equivalence is proved. It is shown that in case of pure curved space-time models without Cartan torsion no specific additional interaction terms because of non-flat geometry arise. The conformal symmetry of a massless generally covariant equation is demonstrated explicitly. A canonical tensor of energy-momentum  $T_{\beta\alpha}$  is constructed, its conservation law happens to involve the Riemann curvature tensor. Within the framework of known ambiguity of any energy-momentum tensor, a new tensor  $\bar{T}_{\beta\alpha}$  is suggested to be used, which obeys a common conservation law.

## References

1. Pauli W., Fierz M. Uber relativistische Feldgleichungen von Teilchen mit beliebigem Spin im electromagnetischen Feld. // *Helv. Phys. Acta.* 1939, Bd. 12, S. 297-300.
2. Fierz V., Pauli W. On relativistic wave equations for particles of arbitrary spin in an electromagnetic field. // *Proc. Roy. Soc. London.* 1939, Vol. A173, P. 211-232.
3. Dirac P. Relativistic wave equations. *Proc. Roy. Soc. (London).* A. 1936. Vol. 155. P. 447-459.
4. H.J. Bhabha. Relativistic wave equations for elementary particles. *Rev. Mod. Phys.* 1945. Vol. 17. P. 200-216; 1949. vol. 21. P. 451; On the postulational basis of the theory of elementary particles. *Rev. Mod. Phys.* 1949. Vol. 21. P. 451-462.
5. Harish-Chandra. *Phys. Rev.* 1947. Vol. 71. P. 793; *Proc. Roy. Soc. (London).* 1947. Vol. 192A. P. 195.
6. H. Umezawa. *Quantum field theory.* Amsterdam. 1956.
7. E.M. Corson. *Introduction to tensors, spinors and relativistic wave equations.* Blackie. London. 1953.
8. I.M. Gel'fand, R.A. Minlos, and Z.Ya. Shapiro. *Linear representations of the rotation and Lorentz groups and their applications.* Pergamon. New York. 1963.
9. M.A. Naimark. *Linear representations of the Lorentz group.* Pergamon. New York, 1964.
10. Krajcik R., Nieti M.M. Bhabha first-order wave equations. *Phys. Rev. D.* 1974. Vol. 10. P. 4049-4062; *Phys. Rev. D.* 1975. Vol. 11. P. 4042-1458; *Phys. Rev. D.* 1975. Vol. 11. P. 4059-1471; *Phys. Rev. D.* 1976. Vol. 13. P. 924-941; *Phys. Rev. D.* 1976. Vol. 14. P. 418-436; *Phys. Rev. D.* 1977. Vol. 15. P. 433-444; *Phys. Rev. D.* 1977. Vol. 15. P. 445-452.
11. A.A. Bogush, L.G. Moroz. *Introduction to theory of classical fields.* Minsk, 1968. 385 pages (in Russian).
12. F.I. Fedorov. *The Lorentz group.* Moskow, 1989. 384 pages (in Russian ).
13. V.L. Ginzburg. *On theory of excited spin states of elementary particles.* *JETP.* 1943. Vol. 13. P. 33-58 (in Russian).
14. E.S. Fradkin. *On theory of particles with higher spins.* *JETP.* 1950. Vol 20. P. 27-38 (in Russian).
15. Petras M. *A note to Bhabha's equation for a particle with maximum spin 3/2.* *Czech. J. Phys.* 1955. Vol. 5. No 3. P. 418-419.
16. V.Ya. Fainberg. *On interaction theory of higher spin particles with electromagnetic and meson fields.* *Trudy FIAN SSSR.* 1955. Vol. 6. P. 269-332 (in Russian).
17. I. Ulehla *Anomalous equations for particles with spin 1/2.* *JETP.* 1957. Vol. 33. P. 473-477 (in Russian).
18. Formanek J. *On the Ulehla-Petras wave equation.* *Czech. J. Phys. B.* 1955. Vol. 25. No 8. P. 545-553.
19. S.I. Lobko. *On theory particles with variable spin 1/2 -3/2 and with two rest masses.* *Kandid. Dissertation.* Minsk, 1965 (in Russian).
20. F.I. Fedorov, V.A. Pletushev. *Wave equations with repeated representations of the Lorentz group. Integer spin.* *Vesti AN BSSR. Ser. fiz.-mat. nauk.* 1969. No 6. P. 81-88 (in Russian).
21. A.Z. Capri. *Non-uniqueness of the spin 1/2 equation.* *Phys. Rev.* 1969. Vol. 178. No 5. P. 1811-1815.
22. A.Z. Capri. *First-order wave equations for half-odd-integral spin.* *Phys. Rev.* 1969. Vol. 178. No 5. P. 2427-2433.
23. A.Z. Capri. *First-order wave equations for multi-mass fermions.* *Nuovo Cimento. B.* 1969. Vol. 64. No 1. P. 151-158.
24. F.I. Fedorov, V.A. Pletushev. *Wave equations with repeated representations of the Lorentz group. Half-integer spin.* *Vesti AN BSSR. Ser. fiz.-mat. nauk.* 1970. No 3. P. 78-83 (in Russian).
25. V.A. Pletushev, F.I. Fedorov. *Wave equations with repeated representations of the Lorentz group for a spin 0 particle.* *Vesti AN BSSR. Ser. fiz.-mat. nauk.* 1970. No 2. P. 79-85 (in Russian).
26. V.A. Pletushev, F.I. Fedorov. *Wave equations with repeated representations of the Lorentz group for a spin 1 particle.* *Vesti AN BSSR. Ser. fiz.-mat. nauk.* 1970, No 3. P. 84-92 (in Russian).
27. A. Shamaly, A.Z. Capri. *First-order wave equations for integral spin.* *Nuovo Cimento. B.* 1971. Vol. 2. No 2. P. 235-253.
28. A.Z. Capri. *Electromagnetic properties of a new spin-1/2 field.* *Progr. Theor. Phys.* 1972. Vol. 48. No 4. P. 1364-1374.
29. A. Shamaly, A.Z. Capri. *Unified theories for massive spin 1 fields.* *Can. J. Phys.* 1973. Vol. 51. No 14. P. 1467-1470.
30. M.A.K. Khalil. *Properties of a 20-component spin 1/2 relativistic wave equation.* *Phys. Rev. D.* 1977. Vol. 15. No 6. P. 1532-1539.

31. M.A.K. Khalil. *Barnacle equivalence structure in relativistic wave equation*. Progr. Theor. Phys. 1978. Vol. 60. No 5. P. 1559-1579.
32. M.A.K. Khalil. *An equivalence of relativistic field equations*. Nuovo Cimento. A. 1978. Vol. 45. No 3. P. 389-404.
33. M.A.K. Khalil. *Reducible relativistic wave equations*. J. Phys. A.: Math. and Gen. 1979. Vol. 12. No 5. P. 649-663.
34. A.A. Bogush, V.V. Kisel. *Equations with repeated representations of the Lorentz group and Pauli interaction*. Vesti AN BSSR. Ser. fiz.-mat. nauk. 1979. No 3. P. 61-65 (in Russian).
35. A.A. Bogush, V.V. Kisel, M.I. Levchuk, L.G. Moroz. *On description of polarizability of scalar particles in the theory of relativistic wave equations*. In: Covariant methods in theoretical physics. Elementary particle physics and relativity theory. Institute of Physics, Academy of sciences of Belarus. Minsk. 1981. P. 81-90 (in Russian).
36. V.V. Kisel. *Electric polarizability of a spin 1 particle in the theory of relativistic wave equations*. Vesti AN BSSR. Ser. fiz.-mat. nauk. 1982. No 3. P. 73-78 (in Russian).
37. A.A. Bogush, V.V. Kisel. *Description of a free particle by different wave equations*. Doklady AN BSSR. 1984. Vol. 28. No 8. P. 702-705 (in Russian).
38. A.A. Bogush, V.V. Kisel. *Equation for a spin 1/2 particle with anomalous magnetic momentum*. // Izvestiya VUZov. Fizika. 1984. No 1. P. 23-27 (in Russian).
39. A.A. Bogush, V.V. Kisel, F.I. Fedorov. *On interpretation of supplementary components of wave functions in presence of electromagnetic interaction*. Doklady AN BSSR. 1984. Vol. 277. No 2. P. 343-346 (in Russian).
40. Lunardi J.T., Pimentel B.M., Teixeira R.G. *Duffin-Kemmer-Petiau equation in Riemannian space-times*. – 14 pages. gr-qc/9909033.
41. Lunardi J. T., Pimentel B.M., Teixeira R. G. and Valverde J.S. 2000. *Remarks on Duffin-Kemmer-Petiau theory and gauge invariance*. Phys. Lett. A. 268. 165-73.
42. Fainberg V. Ya. and Pimentel B.M. *Duffin-Kemmer-Petiau and Klein-Gordon-Fock equations for electromagnetic, Yang-Mills and external gravitational field interactions: proof of equivalence*. Phys. Lett. 2000. A. Vol. 271. P. 16-25.
43. Fainberg V. Ya. and Pimentel B.M. *On equivalence Duffin-Kemmer-Petiau and Klein-Gordon equations*. Braz. J. Phys. 2000. Vol. 30. P. 275-81.
44. Fainberg V. Ya. and Pimentel B.M. *Equivalence Duffin-Kemmer-Petiau and Klein-Gordon-Fock equations*. Theor. Math. Phys. 2000. Vol. 124. P. 1234-49.
45. de Montigny M., Khanna F.C., Santana A. E., Santos E.S. and Vianna J.D.M. *Galilean covariance and the Duffin-Kemmer-Petiau equation*. J.Phys. A.: Math. Gen. 2000. Vol. 33. L273-8.
46. Tetrode H. *Allgemein relativistische Quanten theorie des Elektrons*. Z. Phys. 1928, Bd 50, S. 336.
47. Weyl H. *Gravitation and the electron*. Proc. Nat. Acad. Sci. Amer. – 1929. – Vol. 15. – P. 323-334; *Gravitation and the electron*. Rice Inst. Pamphlet. – 1929. – Vol. 16. – P. 280-295; *Elektron und Gravitation*. Z. Phys. – 1929. – Bd. 56. – S. 330-352; *A remark on the coupling of gravitation and electron*. Actas Akad. Nat. Ciencias Exactas. Fis. y natur. Lima. – 1948. – Vol. 11. – P. 1-17.
48. Fock V., Ivanenko D. *Über eine mögliche geometrische Deutung der relativistischen Quantentheorie*. Z. Phys. – 1929. – Bd. 54. – S. 798-802; *Géometrie quantique linéaire et déplacement parallèle*. Compt. Rend. Acad. Sci. Paris. – 1929. – Vol. 188. – P. 1470-1472.
49. Fock V. *Geometrisierung der Diracschen Theorie des Elektrons*. Z. Phys. – 1929. – Bd. 57, n. 3-4. – S. 261-277; *Sur les équations de Dirac dans la théorie de relativité générale*. Compt. Rend. Acad. Sci. Paris. – 1929. – Vol. 189. – P. 25-28.
50. Schouten J.A. *Dirac equations in general relativity*. J. Math. and Phys. – 1931. – Vol. 10. – P. 239-271; P. 272-283.
51. Schrödinger E. *Sur la théorie relativiste de l'électron et l'interprétation de la mécanique quantique*. Ann. Inst. H. Poincaré. – 1932. – Vol. 2. – P. 269-310; *Dirac'sches Elektron im Schwerfeld*. Sitz. Ber. Preuss. Akad. Wiss. Berlin. Phys.-Math Kl. – 1932. – S. 105-128.
52. Einstein A., Mayer W. *Semivektoren und Spinoren*. Sitz. Ber. Preuss. Akad. Wiss. Berlin. Phys.-Math. Kl. – 1932. – S. 522-550; *Die Diracgleichungen für Semivektoren*. Proc. Akad. Wet. (Amsterdam). – 1933. – Bd. 36. – S. 497-516; *Spaltung der Natürlichsten Feldgleichungen für Semi-Vektoren in Spinor-Gleichungen von Diracschen Typus*. Proc. Akad. Wet. (Amsterdam). – 1933. – Bd. 36. – S. 615-619.
53. Bargmann V. *Über der Zusammenhang zwischen Semivektoren und Spinoren und die Reduktion der Diracgleichungen für Semivektoren*. Helv. Phys. Acta. – 1933. – Bd. 7. – S. 57.



54. Schouten J.A. *Zur generellen Feldtheorie, Semi-Vektoren und Spin-raum.* Z. Phys. – 1933. – Bd. 84. – S. 92-111.
55. Infeld L., van der Waerden B.L. *Die Wellengleichungen des Elektrons in der allgemeinen Relativitätstheorie.* Sitz. Ber. Preuss. Akad. Wiss. Berlin. Phys.-Math. Kl. – 1933. – Bd. 9. – S. 380-401.
56. Infeld L. *Dirac's equation in general relativity theory.* Acta Phys. Polon. – 1934. – Vol. 3, n. 3. – P. 1.
57. Dirac P.A.M. *em Wave equations in conformal space.* Ann. Math. – 1936. – Vol. 37. – P. 429-442.
58. Yamamoto H. *On equations for the Dirac electron in general relativity.* Japan. J. Phys. – 1936. – Vol. 11. – P. 35.
59. Proca A. *Sur la théorie ondulatoire des electrons positifs et négatifs.* J. Phys. et Rad. – 1936. – Vol. 7. – P. 347.
60. Benedictus W. *Les équations de Dirac dans un espace à métrique riemannienne.* Compt. Rend. Acad. Sci. Paris. – 1938. – Vol. 206. – P. 1951.
61. Cartan E. *Leçons sur la théorie des spineurs.* Actualités Sci. – 1938.
62. Pauli W. *Über die Invarianz der Dirac'schen Wellengleichungen gegenüber Ähnlichkeit stransformationen des Linienelementes im Fall verschwindender Ruhmasse.* Helv. Phys. Acta. – 1940. – Bd. 13. – S. 204-208.
63. Bade W.L., Jehle H. *An introduction to spinors.* Rev. Mod. Phys. – 1953. – Vol. 25. – P. 714-728.
64. Brill D.R., Wheeler J.A. *Interaction of neutrinos and gravitational fields.* Rev. Mod. Phys. – 1957. – Vol. 29, n. 3. – P. 465-479.
65. Bergmann P.G. *Two-component spinors in general relativity.* Phys. Rev. – 1957. – Vol. 107, n. 2. – P. 624-629.
66. Fletcher J.G. *Dirac matrices in Rimannian space.* Nuovo Cim. – 1958. – Vol. 8, n. 3. – P. 451-458.
67. Buchdahl H.A. *On extended conformal transformations of spinors and spinor equations.* Nuovo Cim. – 1959. – Vol. 11. – P. 496-506.
68. Namyslowski J. *The Dirac equation in general relativity in the vierbein formalism.* Acta Phys. Polon. – 1961. – Vol. 20, n. 11. – P. 927-936; *Symmetrical form of Dirac matrices in general relativity.* Acta Phys. Polon. – 1963. – Vol. 23, n. 6. – P. 673-684.
69. Peres A. *Spinor fields in generally covariant theories.* Nuovo Cim. Suppl. – 1962. – Vol. 24, n. 2. – P. 389-452.
70. Lichnerowicz A. *Champ de Dirac, champ du neutrino et transformations C,P,T sur un espace-temps curve.* Ann. Inst. Henri Poincaré. A. 1964, Vol. 1, n. 3, P. 233-290.
71. Ogivetskiy V.I., Polubarinov I.V. *On spinors in gravity theory* (in Russian). JETP. 1965. Vol 48, n. 6. P. 1625-1636.
72. Pagels H. *Spin and Gravitation.* Ann. Phys. (N.Y.)– 1965. – Vol. 31, n. 1. – P. 64-87.
73. Brill D.R., Cohen J.M. *Cartan frames and general relativistic Dirac equation.* J. Math. Phys. – 1966. – Vol. 7, n. 2. – P. 238-245.
74. Cap F., Majerotto W., Raab W., Unteregger P. *Spinor calculus in Riemannian manifolds.* Fortschr. der Physik. – 1966. – Bd. 14, n. 3. – S. 205-233.
75. Von D. Kramer, Stephani H. *Bispinorfelder im Riemannschen Raum.* Acta Phys. Polonica. – 1966. – Vol. 29, n. 3. – P. 379-386.
76. Luehr C.P., Rosenbaum M. *Spinor connections in general relativity.* J. Math. Phys. – 1974. – Vol. 15, n. 7. – P. 1120-1137.
77. Maia M.D. *Conformal spinor fields in general relativity.* J. Math. Phys. – 1974. – Vol. 15, n. 4. – P. 420-425.
78. Brumâ C. *lem A mode of constructing the Dirac matrices in gravitational field.* Rev. Roum. Phys. – 1986. – Tome 31, n. 8. – P. 753-763; *On Dirac matrices in gravitational field.* Rev. Roum. Phys. – 1987. – Tome 32, n. 4. – P. 375-382.
79. Fariborz A.H., McKeon D.G.C. *Spinors in Weyl geometry.* Class. Quant. Grav. – 1997. – Vol. 14. – P. 2517-2525. – hep-th/9607056.
80. Manoelito M. de Souza. *The Lorentz Dirac equation and the structure of spacetime.* Braz. J. Phys. – 1998. – Vol. 28. – P. 250-256. – hep-th/9505169.